

Periodic solutions of pendulum-like Hamiltonian systems in the plane

Alessandro Fonda, Rodica Toader

Dedicated to Jean Mawhin, in the occasion of his 70th birthday

ABSTRACT. By the use of a generalized version of the Poincaré–Birkhoff fixed point theorem, we prove the existence of at least two periodic solutions for a class of Hamiltonian systems in the plane, having in mind the forced pendulum equation as a particular case. Our approach is closely related to the one used by Franks in [14]. We thus provide a new proof of a theorem by Mawhin and Willem [26], originally obtained by the use of variational methods.

1 Introduction

For many years, the periodically forced pendulum equation

$$\ddot{x} + a \sin x = e(t) \tag{1}$$

has been a fruitful source of mathematical ideas. We refer to [24] for a comprehensive review on this subject.

In 1922, Hamel [17] proved that, if $e(t)$ is T -periodic and has mean value equal to zero, i.e.,

$$\int_0^T e(t) dt = 0, \tag{2}$$

then equation (1) has at least one T -periodic solution. This result, forgotten for a long time, was rediscovered independently by Dancer [8] and Willem [36].

In 1984, Mawhin and Willem [26] proved that, if the above assumption (2) holds, equation (1) has indeed at least two geometrically distinct T -periodic solutions (i.e., solutions not differing by a multiple of 2π). Their proof uses variational methods; the first solution is obtained by minimizing the action functional, as already observed in [8, 17, 36]. (In the autonomous case, this solution corresponds to the unstable equilibrium of the pendulum.) Clearly, by the periodicity in x , this means that there are infinitely many minimum points, differing from each other by a multiple of 2π . So, considering two of them and applying a mountain pass argument, a critical point of saddle type is obtained, which corresponds to a solution which differs from all the previous

ones. (In the autonomous case, this new solution corresponds to the stable equilibrium.) The result by Mawhin and Willem, as already noticed in [26], continues to hold if the restoring force $a \sin x$ is replaced by a Carathéodory function of the type $g(t, x)$, with a 2π -periodic primitive $G(t, x) = \int_0^x g(t, s) ds$.

In 1988, Franks [14] (see also [15]) provided a new proof of the existence of at least two T -periodic solutions, by the use of a variant of the Poincaré–Birkhoff fixed point theorem on a cylindrical annulus. His idea is to consider the flow generated by the equivalent first order system on the cylinder obtained by identifying $x = 0$ with $x = 2\pi$, observing that, far from the origin, a twist type property holds on the upper and lower sections of the cylinder.

Recently, in [3] and [2], similar results were obtained for the so-called “relativistic pendulum” equation

$$\frac{d}{dt} \frac{\dot{x}}{\sqrt{1 - (\dot{x}/c)^2}} + a \sin x = e(t),$$

by the use of variational methods combined with topological and nonsmooth analysis techniques. Under the same assumption (2), the existence of at least two geometrically distinct T -periodic solutions was proved in [2]. On the other hand, in [29], the existence of two T -periodic solutions was proved for the equation

$$\frac{d}{dt} \frac{\dot{x}}{\sqrt{1 + (\dot{x}/\gamma)^2}} + a \sin x = e(t),$$

involving a “mean curvature” type operator, where, besides (2), a boundedness condition on $E(t) = \int_0^t e(\tau) d\tau$ was assumed, i.e., that $\|E\|_\infty < \gamma$. In this case, however, the notion of solution has been extended to the class of functions of bounded variation, so that a solution may exhibit some points of discontinuity.

The pendulum equation has been extended in different directions to systems of differential equations of the type

$$\ddot{x} + \nabla V(t, x) = 0,$$

with $x(t) \in \mathbb{R}^N$, or to Hamiltonian systems like

$$J\dot{u} = \nabla H(t, u), \tag{3}$$

with $u(t) \in \mathbb{R}^{2N}$, where J is the standard $2N \times 2N$ symplectic matrix, namely

$$J = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}.$$

See, e.g., [5, 6, 7, 10, 11, 18, 22, 27, 30, 32, 33, 34], where multiplicity results for the periodic problem were proved by variational methods. Following Conley and Zehnder [6, 7], these results are somewhat related to a conjecture by Arnold [1], concerning a possible version of the Poincaré–Birkhoff fixed point theorem in higher dimensions. Recently, some of these results have been extended in [4, 25] to systems of relativistic-type pendulums.

Let us also recall that, under the same assumption (2), the existence of infinitely many quasi-periodic solutions for the pendulum equation has been proved in [20, 37] by the use of the Kolmogorov-Arnold-Moser theory.

The aim of this paper is to generalize the result obtained by Mawhin and Willem for the pendulum equation to a planar Hamiltonian system of the type (3), by the use of a generalized version of the Poincaré–Birkhoff fixed point theorem on a planar annulus (see [9, 31]). Writing $u = (x, y)$ we will assume the Hamiltonian function $H(t, x, y)$ to have a linear growth, to be 2π -periodic in x , and to satisfy some kind of monotonicity condition in y , for $|y|$ large enough. More precisely, we will ask the function $y \mapsto H(t, x, y)$ to be strictly increasing when y is large and positive, and strictly decreasing for y large and negative.

The idea of the proof is to transform the Hamiltonian system into a new one, where x plays the role of an angle and y is related to the radial displacement. For a sufficiently large positive number α , we restrict our analysis to the half plane

$$\Omega_\alpha = \{(x, y) \in \mathbb{R}^2 : y > -\alpha\},$$

and, by the transformation

$$\varphi_\alpha(x, y) = \left(\sqrt{2(y + \alpha)} \cos x, -\sqrt{2(y + \alpha)} \sin x \right),$$

we obtain a new Hamiltonian system to which the Poincaré–Birkhoff theorem can be applied.

Our method applies to the relativistic pendulum equation and to equations involving the curvature operator, as well. It also provides, under suitable assumptions, the existence and multiplicity of the so-called “periodic solutions of the second kind”, i.e., those satisfying, for some integer k ,

$$u(t + T) = u(t) + (2\pi k, 0), \quad \text{for every } t \in \mathbb{R}.$$

Such a problem has already been faced for pendulum-like equations in [23]. We will also consider the existence of “subharmonic solutions of the second kind”, defined as above, when T is replaced by an integer multiple of T .

2 The main results

We consider the planar Hamiltonian system

$$J\dot{u} = \nabla H(t, u) \tag{4}$$

where $H : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a Carathéodory function, T -periodic in the first variable t , and such that $\nabla H : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, the gradient with respect to the second variable u , is locally Lipschitz continuous in u . More precisely,

(i) $H(\cdot, u)$ is measurable, for every $u \in \mathbb{R}^2$;

- (ii) $H(t, \cdot)$ is differentiable, for almost every $t \in \mathbb{R}$;
- (iii) for every compact set $K \subseteq \mathbb{R}^2$, there exists a function $\ell_K \in L^1(0, T)$ such that, for almost every $t \in [0, T]$ and for every $u_1, u_2 \in K$,

$$|\nabla H(t, u_1) - \nabla H(t, u_2)| \leq \ell_K(t)|u_1 - u_2|.$$

We look for T -periodic solutions of system (4). To this aim, we introduce the following three assumptions. The first one says that $\nabla H(t, u)$ has an at most linear growth in u .

Assumption A1. *There are a constant $c_1 \geq 0$ and a function $c_2 \in L^1(0, T)$ such that*

$$|\nabla H(t, u)| \leq c_1|u| + c_2(t), \quad \text{for a.e. } t \in [0, T] \text{ and every } u \in \mathbb{R}^2.$$

In order to state the next two assumptions, it is more convenient to write system (4) in its equivalent form

$$\dot{x} = \frac{\partial H}{\partial y}(t, x, y), \quad \dot{y} = -\frac{\partial H}{\partial x}(t, x, y). \quad (5)$$

We now ask that $H(t, x, y)$ be 2π -periodic in x .

Assumption A2. *The function H satisfies*

$$H(t, x + 2\pi, y) = H(t, x, y), \quad \text{for a.e. } t \in [0, T] \text{ and every } (x, y) \in \mathbb{R}^2.$$

In the third assumption we impose a sign condition on $\frac{\partial H}{\partial y}(t, x, y)$ when $|y|$ is large enough.

Assumption A3. *There is a constant $d > 0$ such that*

$$|y| \geq d \quad \Rightarrow \quad \text{sgn}(y) \frac{\partial H}{\partial y}(t, x, y) > 0,$$

for almost every $t \in [0, T]$ and every $x \in [0, 2\pi]$.

We can now state our main result. We say that two solutions are *geometrically distinct* if they do not differ by a multiple of 2π . In the sequel, when quoting multiplicity results, we always refer to geometrically distinct solutions, even if not explicitly stated.

Theorem 1 *If Assumptions A1, A2, and A3 hold, then system (4) has at least two geometrically distinct T -periodic solutions.*

As an example, let us consider the forced pendulum equation

$$\ddot{x} + a \sin x = e(t), \quad (6)$$

where $e : \mathbb{R} \rightarrow \mathbb{R}$ is a T -periodic, locally integrable function, satisfying the zero mean condition (2). Setting $E(t) = \int_0^t e(\tau) d\tau$, we can write the equivalent system

$$\dot{x} = y + E(t), \quad \dot{y} = -a \sin x.$$

We thus have a Hamiltonian system, with

$$H(t, x, y) = \frac{1}{2} y^2 + E(t) y - a \cos x,$$

which clearly satisfies Assumptions A1, A2 and A3. So, Theorem 1 guarantees the existence of at least two T -periodic solutions to equation (6), a result first proved by Mawhin and Willem [26].

The same will be true for an equation like

$$\frac{d}{dt} \phi(\dot{x}) + g(t, x) = e(t). \quad (7)$$

Here, $\phi :]-c, c[\rightarrow \mathbb{R}$, with $c \in]0, +\infty[$, is a diffeomorphism for which

$$\inf\{\phi'(s) : s \in]-c, c[\} > 0. \quad (8)$$

Moreover, the function $g(t, x)$ is Carathéodory, T -periodic in t , locally Lipschitz continuous and 2π -periodic in x , and such that

$$\int_0^{2\pi} g(t, x) dx = 0, \quad (9)$$

while the function $e : \mathbb{R} \rightarrow \mathbb{R}$ is T -periodic, locally integrable, and satisfying (2).

Writing the equivalent system

$$\dot{x} = \phi^{-1}(y + E(t)), \quad \dot{y} = -g(t, x), \quad (10)$$

it is easily seen that Assumptions A1, A2 and A3 are satisfied, so that Theorem 1 guarantees the existence of at least two T -periodic solutions to equation (7). We thus recover a recent result by Bereanu and Torres [2]. Notice however that, in [2], the function $g(t, x)$ was not assumed to be locally Lipschitz continuous in x . As an example, we could have

$$\phi(s) = \frac{s}{\sqrt{1 - (s/c)^2}}, \quad (11)$$

leading to the so-called “relativistic pendulum”, see [2, 3].

In Section 5 we will see how to generalize this situation so to consider a diffeomorphism $\phi :]-c, c[\rightarrow]-\gamma, \gamma[$, satisfying (8), with $c, \gamma \in]0, +\infty[$, thus including also the case of the mean curvature operator.

Let us also recall a related result proved in [5, 10, 11, 32] for a Hamiltonian system in \mathbb{R}^{2N} like

$$J\dot{u} = \mathbb{A}u + \nabla\mathcal{H}(t, u), \quad (12)$$

where \mathbb{A} is a symmetric matrix such that the only T -periodic solutions of $J\dot{u} = \mathbb{A}u$ are the constants in $\ker \mathbb{A}$. It was proved that, if $\mathcal{H}(t, \cdot)$ is T -periodic along the directions of $\ker \mathbb{A}$, and has bounded gradient and bounded hessian matrix in u , then system (12) has at least $\dim(\ker \mathbb{A}) + 1$ periodic solutions of period T . If we now assume that $N = 1$ and take

$$\mathbb{A} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix},$$

then system (12) can be written as

$$\dot{x} = y + \frac{\partial\mathcal{H}}{\partial y}(t, x, y), \quad \dot{y} = -\frac{\partial\mathcal{H}}{\partial x}(t, x, y).$$

If $\nabla\mathcal{H}$ is assumed to be bounded, one has that our Assumptions A1 and A3 are satisfied. Moreover, the periodicity along $\ker \mathbb{A}$ implies our Assumption A2. Hence, in this case, Theorem 1 provides the existence of two T -periodic solutions. Notice that, in order to apply the results in the above quoted papers, we would need T not to be an integer multiple 2π , so to ensure that the only T -periodic solutions of $J\dot{u} = \mathbb{A}u$ be the constants. This condition is no more necessary here.

3 Proof of Theorem 1

Let us first state the version of the Poincaré–Birkhoff theorem [9, 31] which we will use in the proof of our main result (see also [12] for a review on this theorem). In the statement, given a simple closed curve Γ in \mathbb{R}^2 , we denote by $I(\Gamma)$ the open domain bounded by Γ , and by $\bar{I}(\Gamma)$ its closure.

Theorem 2 *Let $\mathcal{A} = \bar{I}(\Gamma_2) \setminus I(\Gamma_1)$, with $0 \in I(\Gamma_1)$, be an annular region bounded by two curves Γ_1 and Γ_2 , which are strictly star-shaped with respect to the origin. Let $\Psi : \bar{I}(\Gamma_2) \rightarrow \Psi(\bar{I}(\Gamma_2))$ be an area-preserving homeomorphism such that $0 \in \Psi(I(\Gamma_1))$. On the universal covering space $\{(\theta, \rho) : \theta \in \mathbb{R}, \rho > 0\}$, with the standard covering projection $\Pi : (\theta, \rho) \mapsto (\rho \cos \theta, \rho \sin \theta)$, consider a lifting of $\Psi|_{\mathcal{A}}$ of the form*

$$h(\theta, \rho) = (\theta + \gamma(\theta, \rho), \eta(\theta, \rho)),$$

where $\gamma(\theta, \rho)$ and $\eta(\theta, \rho)$ are continuous and 2π -periodic in their first variable. Correspondingly, for $\tilde{\Gamma}_1 = \Pi^{-1}(\Gamma_1)$ and $\tilde{\Gamma}_2 = \Pi^{-1}(\Gamma_2)$, assume the twist condition

$$\gamma(\theta, \rho) > 0 \text{ on } \tilde{\Gamma}_1, \quad \gamma(\theta, \rho) < 0 \text{ on } \tilde{\Gamma}_2.$$

Then, Ψ has two fixed points v_1, v_2 in the interior of \mathcal{A} , such that

$$\gamma(\Pi^{-1}(v_1)) = \gamma(\Pi^{-1}(v_2)) = 0.$$

We now start the proof of Theorem 1. For any $u_0 \in \mathbb{R}^2$, let us denote by $u(t; t_0, u_0)$ the solution of the Cauchy problem

$$\begin{cases} J\dot{u} = \nabla H(t, u) \\ u(t_0) = u_0, \end{cases} \quad (13)$$

which is unique and globally defined, by the assumptions made on H . Writing $u_0 = (x_0, y_0)$, and correspondingly

$$u(t; t_0, u_0) = (x(t; t_0, x_0, y_0), y(t; t_0, x_0, y_0)),$$

we see that the periodicity Assumption A2 implies that

$$\begin{aligned} x(t; t_0, x_0 + 2\pi, y_0) &= x(t; t_0, x_0, y_0) + 2\pi, \\ y(t; t_0, x_0 + 2\pi, y_0) &= y(t; t_0, x_0, y_0). \end{aligned} \quad (14)$$

Moreover, as a consequence of the linear growth Assumption A1 and the periodicity Assumption A2, we have that

$$\left| \frac{\partial H}{\partial x}(t, x, y) \right| \leq c_1|y| + \tilde{c}_2(t), \quad (15)$$

with $\tilde{c}_2(t) = 2\pi c_1 + c_2(t)$, for almost every $t \in \mathbb{R}$ and every $(x, y) \in \mathbb{R}^2$. The following proposition is a rather classical application of the Gronwall inequality, once (15) is known.

Proposition 1 *One has*

$$\lim_{y_0 \rightarrow +\infty} y(t; t_0, x_0, y_0) = +\infty, \quad \lim_{y_0 \rightarrow -\infty} y(t; t_0, x_0, y_0) = -\infty,$$

uniformly for every $t, t_0 \in [0, T]$ and $x_0 \in [0, 2\pi]$.

Let $d > 0$ be the constant introduced in Assumption A3. By Proposition 1 there is a $D \geq d$ such that

$$\begin{aligned} y_0 \leq -D &\Rightarrow y(t; 0, x_0, y_0) \leq -d, \\ y_0 \geq D &\Rightarrow y(t; 0, x_0, y_0) \geq d, \end{aligned} \quad (16)$$

for every $t \in [0, T]$ and $x_0 \in [0, 2\pi]$. Moreover, there is a $\beta > D$ such that

$$y_0 \geq -D \Rightarrow y(t; 0, x_0, y_0) \geq -\beta, \quad \text{for every } t \in [0, T] \text{ and } x_0 \in [0, 2\pi]. \quad (17)$$

We will now transform system (4) into a new system, for which it will be possible to apply the Poincaré–Birkhoff fixed point theorem. Let us fix an $\alpha > \beta$. Set

$$\Omega_\alpha = \{(x, y) \in \mathbb{R}^2 : y > -\alpha\},$$

and let $\varphi_\alpha : \Omega_\alpha \rightarrow \mathbb{R}^2 \setminus \{(0, 0)\}$ be defined as

$$\varphi_\alpha(x, y) = \left(\sqrt{2(y + \alpha)} \cos x, -\sqrt{2(y + \alpha)} \sin x \right).$$

Notice that the determinant of the Jacobian matrix $\varphi'_\alpha(x, y)$ is equal to 1, for every $(x, y) \in \Omega_\alpha$. Setting

$$\widehat{\Omega}_\alpha = \{(x, y) \in \mathbb{R}^2 : 0 < x < 2\pi, y > -\alpha\},$$

we thus have that the function

$$\widehat{\varphi}_\alpha : \widehat{\Omega}_\alpha \rightarrow \mathbb{R}^2 \setminus \{(\xi, \eta) \in \mathbb{R}^2 : \xi \geq 0, \eta = 0\},$$

with $\widehat{\varphi}_\alpha(u) = \varphi_\alpha(u)$, is a symplectic diffeomorphism, i.e.,

$$\widehat{\varphi}'_\alpha(u)^T J \widehat{\varphi}'_\alpha(u) = J, \quad (18)$$

for every $u \in \widehat{\Omega}_\alpha$, cf. [16]. (Here \mathbb{M}^T denotes the transposed of a matrix \mathbb{M} .) Then, using (18), one easily sees that the function $\widehat{\varphi}_\alpha$ transforms the Hamiltonian system (4) on $\widehat{\Omega}_\alpha$ into the Hamiltonian system

$$J\dot{v} = \nabla L_\alpha(t, v), \quad (19)$$

on $\mathbb{R}^2 \setminus \{(\xi, \eta) \in \mathbb{R}^2 : \xi \geq 0, \eta = 0\}$, where

$$L_\alpha(t, v) = H(t, \widehat{\varphi}_\alpha^{-1}(v)).$$

(As usual, we denote by ∇L_α the gradient with respect to the second variable.) By the periodicity Assumption A2, we can now extend the new Hamiltonian function L_α to $\mathbb{R}^2 \setminus \{(0, 0)\}$, preserving the regularity assumptions. We will still denote by $L_\alpha(t, v)$ such an extension. Notice that, if $u = (x, y)$ refers to the original system (4), then, for the new system (19), x plays the role of the angular displacement.

We now need to modify (19) in order to apply Theorem 2. Consider a C^2 -function $\chi : [0, +\infty[\rightarrow [0, +\infty[$, such that

$$\chi(r) = \begin{cases} 1 & \text{if } r \geq \sqrt{2(\alpha - \beta)}, \\ 0 & \text{if } r \leq \frac{1}{2}\sqrt{2(\alpha - \beta)}. \end{cases}$$

Let $\widetilde{L}_\alpha : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined as

$$\widetilde{L}_\alpha(t, v) = \begin{cases} \chi(|v|)L_\alpha(t, v) & \text{if } v \neq 0, \\ 0 & \text{if } v = 0. \end{cases}$$

This is a Carathéodory function, T -periodic in the first variable t , and such that $\nabla \widetilde{L}_\alpha$, the gradient with respect to the second variable v , is locally Lipschitz continuous and has an at most linear growth in v . Since \widetilde{L}_α and L_α coincide on the set $\{v \in \mathbb{R}^2 : |v| \geq \sqrt{2(\beta - \alpha)}\}$, we have that the Hamiltonian system

$$J\dot{v} = \nabla \widetilde{L}_\alpha(t, v) \quad (20)$$

still describes the original system (4), as long as $u = (x, y)$ satisfies $y \geq -\beta$. We will denote by $\mathcal{P} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ the Poincaré map associated to this new system.

Let us define the annulus

$$\begin{aligned}\mathcal{A} &= \varphi_\alpha(\overline{\Omega}_{-D} \setminus \Omega_D) \\ &= \overline{B}\left((0, 0), \sqrt{2(D + \alpha)}\right) \setminus B\left((0, 0), \sqrt{2(-D + \alpha)}\right).\end{aligned}$$

Notice that L_α and \tilde{L}_α coincide on \mathcal{A} . We want to apply Theorem 2 to the map \mathcal{P} over the annulus \mathcal{A} . We know that \mathcal{P} is an area-preserving homeomorphism, and since it coincides with the identity on the disk $B\left((0, 0), \frac{1}{2}\sqrt{2(\alpha - \beta)}\right)$, we have that $\mathcal{P}(0, 0) = (0, 0)$.

We now need to check the twist condition. If $|v_0| = \sqrt{2(-D + \alpha)}$, let $x_0 \in [0, 2\pi[$ be such that

$$\varphi_\alpha^{-1}(v_0) = \{(x_0 + 2k\pi, -D) : k \in \mathbb{Z}\}.$$

Then, by (16) and (17),

$$-\beta \leq y(t; 0, x_0, -D) \leq -d, \quad \text{for every } t \in [0, T], \quad (21)$$

and, by Assumption A3, $x(\cdot; 0, x_0, -D)$ is strictly decreasing on $[0, T]$, so we conclude that $x(T; 0, x_0, -D) < x_0$.

Similarly, if $|v_0| = \sqrt{2(D + \alpha)}$, taking $x_0 \in [0, 2\pi[$ for which

$$\varphi_\alpha^{-1}(v_0) = \{(x_0 + 2k\pi, D) : k \in \mathbb{Z}\},$$

we have that $y(t; 0, x_0, D) \geq d$, for every $t \in [0, T]$. Hence, by Assumption A3, $x(\cdot; 0, x_0, D)$ is strictly increasing on $[0, T]$, so that $x(T; 0, x_0, D) > x_0$.

Since, as long as $y \geq -\beta$, we know that x gives the angular displacement for system (20), the twist condition is verified, and Theorem 2 applies, providing the existence of two fixed points $v_0^{(1)}, v_0^{(2)}$ of \mathcal{P} in \mathcal{A} , such that, setting $u_0^{(j)} = (x_0^{(j)}, y_0^{(j)}) = \tilde{\varphi}_\beta^{-1}(v_0^{(j)})$, one has

$$x(T; 0, x_0^{(j)}, y_0^{(j)}) = x_0^{(j)},$$

for $j = 1, 2$. Therefore, $u_0^{(1)}$ and $u_0^{(2)}$ are the starting points of the two T -periodic solutions of system (4) we are looking for. The proof is thus completed.

4 Periodic solutions of the second kind

In this section we study the existence of the so-called ‘‘periodic solutions of the second kind’’, those satisfying, for some integer k ,

$$u(t + T) = u(t) + (2\pi k, 0), \quad \text{for every } t \in \mathbb{R}. \quad (22)$$

These are sometimes called ‘‘running solutions’’. Clearly, if $k = 0$, we recover the usual periodicity. Notice that (22) is equivalent to ask that the function

$$\tilde{u}_1(t) = u(t) - \left(\frac{2\pi k}{T}t, 0\right)$$

be T -periodic. Let us introduce a new assumption.

Assumption A4. *There are two integers k_1, k_2 , with $k_1 \leq k_2$ and two constants $d_1 < d_2$, such that*

$$\begin{aligned} y \leq d_1 &\Rightarrow \frac{\partial H}{\partial y}(t, x, y) < \frac{2\pi k_1}{T}, \\ y \geq d_2 &\Rightarrow \frac{\partial H}{\partial y}(t, x, y) > \frac{2\pi k_2}{T}, \end{aligned}$$

for almost every $t \in [0, T]$ and every $x \in [0, 2\pi]$.

The following theorem generalizes Theorem 1.

Theorem 3 *If Assumptions A1, A2, and A4 hold, then, for every integer $k \in \{k_1, k_1 + 1, \dots, k_2\}$, system (4) has at least two geometrically distinct periodic solutions of the second kind, satisfying (22).*

The proof is a straightforward modification of the proof of Theorem 1. The main difference appears when applying the Poincaré–Birkhoff theorem, in the version provided by Theorem 2. Instead of considering the Poincaré map \mathcal{P} , we have to take the function $\Psi = \mathcal{R}_k \circ \mathcal{P}$, where \mathcal{R}_k denotes the rotation of angle $-2\pi k$, with $k \in \{k_1, k_1 + 1, \dots, k_2\}$. Assumption A4 then provides the necessary twist property.

For a second order differential equation like (7), we have the following.

Corollary 1 *Taking $\phi :]-c, c[\rightarrow \mathbb{R}$ as in Section 2, with $c \in]0, +\infty[$, for any integer k in the interval $]-\frac{T}{2\pi}c, \frac{T}{2\pi}c[$, equation (7) has at least two geometrically distinct periodic solutions of the second kind, satisfying*

$$x(t + T) = x(t) + 2\pi k, \quad \text{for every } t \in \mathbb{R}.$$

Proof It is sufficient to observe that, in this case,

$$\begin{aligned} \lim_{y \rightarrow -\infty} \frac{\partial H}{\partial y}(t, x, y) &= \lim_{y \rightarrow -\infty} \phi^{-1}(y + E(t)) = -c, \\ \lim_{y \rightarrow +\infty} \frac{\partial H}{\partial y}(t, x, y) &= \lim_{y \rightarrow +\infty} \phi^{-1}(y + E(t)) = c, \end{aligned}$$

uniformly for every $t \in [0, T]$ and every $x \in [0, 2\pi]$. If c is finite, we take k_2 to be the largest integer less than $\frac{T}{2\pi}c$; otherwise, if $c = +\infty$, k_2 can be an arbitrary positive integer. Setting $k_1 = -k_2$, we see that Assumption A4 is verified. Theorem 3 then applies. \blacksquare

We could also look for “subharmonic solutions of the second kind”, those satisfying, for some integer k , and for some positive integer m ,

$$u(t + mT) = u(t) + (2\pi k, 0), \quad \text{for every } t \in \mathbb{R}. \quad (23)$$

To this aim, we introduce the following.

Assumption A5. For some positive integer m , there are two integers k_1, k_2 , with $k_1 \leq k_2$ and two constants $d_1 < d_2$, such that

$$\begin{aligned} y \leq d_1 &\Rightarrow \frac{\partial H}{\partial y}(t, x, y) < \frac{2\pi k_1}{mT}, \\ y \geq d_2 &\Rightarrow \frac{\partial H}{\partial y}(t, x, y) > \frac{2\pi k_2}{mT}, \end{aligned}$$

for almost every $t \in [0, T]$ and every $x \in [0, 2\pi]$.

Let us now state the corresponding existence result.

Theorem 4 *If Assumptions A1, A2, and A5 hold, then, for every nonzero integer $k \in \{k_1, k_1 + 1, \dots, k_2\}$, with $|k|$ and m relatively prime, system (4) has at least two geometrically distinct subharmonic solutions of the second kind, satisfying (23), for which there is no other $\ell \in \{1, 2, \dots, m - 1\}$ such that, for some integer j ,*

$$u(t + \ell T) = u(t) + (2\pi j, 0), \quad \text{for every } t \in \mathbb{R}. \quad (24)$$

Proof Let k and m be as in the statement. By Theorem 3, with T replaced by mT , we find two solutions satisfying (23). Let u be one of them. Assume by contradiction that (24) holds, for some integers $\ell \in \{1, 2, \dots, m - 1\}$ and j . Let p and n be positive integers such that $p\ell = nm$. By (23), it has to be

$$u(t + nmT) = u(t) + (2\pi kn, 0), \quad \text{for every } t \in \mathbb{R},$$

and, by (24),

$$u(t + p\ell T) = u(t) + (2\pi jp, 0), \quad \text{for every } t \in \mathbb{R}.$$

Hence, $nk = pj$, so that

$$\frac{k}{m} = \frac{j}{\ell},$$

a contradiction, since $|k|$ and m are relatively prime, and $1 \leq \ell \leq m - 1$. ■

5 Further results

In this section we discuss on possible generalizations of our assumptions and on different applications of our results.

As an alternative to Assumption A3, let us introduce the following.

Assumption A3'. There exists a $\zeta \in L^1(0, T)$ such that

$$\operatorname{sgn}(y) \frac{\partial H}{\partial y}(t, x, y) \geq \zeta(t),$$

for almost every $t \in [0, T]$ and every $(x, y) \in \mathbb{R}^2$, and

$$\int_0^T \limsup_{y \rightarrow -\infty} \left(\sup_{x \in [0, 2\pi]} \frac{\partial H}{\partial y}(t, x, y) \right) dt < 0 < \int_0^T \liminf_{y \rightarrow +\infty} \left(\inf_{x \in [0, 2\pi]} \frac{\partial H}{\partial y}(t, x, y) \right) dt.$$

This is a Landesman-Lazer type of condition. Notice, however, that such kind of condition usually regards the inferior and superior limits as x goes to $\pm\infty$, when dealing with second order scalar equations. Here, since the nonlinearity is periodic in x , and we deal with a more general Hamiltonian system, we propose the above version, with the inferior and superior limits as y goes to $\pm\infty$.

We can then state the following variant of Theorem 1.

Theorem 5 *If Assumptions A1, A2, and A3' hold, then system (4) has at least two geometrically distinct T -periodic solutions.*

Proof Following [13, Lemma 1], if Assumption A3' holds, there are a constant $d \geq 1$ and two functions $\psi_1, \psi_2 \in L^1(0, T)$ such that

$$\begin{aligned} \frac{\partial H}{\partial y}(t, x, y) &\leq \psi_1(t), \text{ for a.e. } t \in [0, T], \text{ every } x \in [0, 2\pi], \text{ and every } y \leq -d, \\ \frac{\partial H}{\partial y}(t, x, y) &\geq \psi_2(t), \text{ for a.e. } t \in [0, T], \text{ every } x \in [0, 2\pi], \text{ and every } y \geq d, \end{aligned}$$

and

$$\int_0^T \psi_1(t) dt < 0 < \int_0^T \psi_2(t) dt.$$

We now follow the lines of the proof of Theorem 1, until we obtain (21), when $y_0 = -D$. At this point, writing for simplicity $x(t)$, $y(t)$ instead of $x(t; 0, x_0, -D)$, $y(t; 0, x_0, -D)$, respectively, we have that

$$x(T; 0, x_0, -D) - x_0 = \int_0^T \frac{\partial H}{\partial y}(t, x(t), y(t)) dt \leq \int_0^T \psi_1(t) dt < 0,$$

so that $x(T; 0, x_0, -D) < x_0$. Similarly, when $y_0 = D$, we get $x(T; 0, x_0, D) > x_0$, and the proof is completed. \blacksquare

Let us now focus our attention on Assumption A1: It has been used in order that the Poincaré map be well defined, and to obtain Proposition 1. To this aim, it would be enough to assume global existence for the solutions of the Cauchy problems (13), cf. [19]. Or else, we could try to follow the proof of Theorem 1 anyway, being careful that the Poincaré map be well defined on the annulus we are interested in, and obtaining the desired estimates anyway.

To show how this can be done, consider again the equation

$$\frac{d}{dt}\phi(\dot{x}) + g(t, x) = e(t), \quad (25)$$

and assume now that $\phi :] - c, c[\rightarrow] - \gamma, \gamma[$ is a diffeomorphism satisfying (8), with $c, \gamma \in]0, +\infty[$, and such that $\phi(0) = 0$. As before, the function $g(t, x)$ is assumed to be Carathéodory, T -periodic in t , locally Lipschitz continuous and 2π -periodic in x , and such that (9) holds. The function $e : \mathbb{R} \rightarrow \mathbb{R}$ is T -periodic, locally integrable, and satisfying (2), so that we can write the equation in its equivalent form (10).

By the above assumptions, there is an $h \in L^1(0, T)$ for which

$$|g(t, x)| \leq h(t), \quad \text{for a.e. } t \in [0, T] \text{ and every } x \in \mathbb{R}.$$

Set, for $\varepsilon > 0$ sufficiently small,

$$\tilde{d} = \|E\|_\infty + \varepsilon, \quad \tilde{D} = \tilde{d} + \|h\|_1, \quad \tilde{\beta} = \tilde{D} + \|h\|_1.$$

Then, for any solution of the corresponding Cauchy problem, since

$$|\dot{y}(t; 0, x_0, y_0)| \leq h(t), \quad \text{for a.e. } t \in [0, T],$$

we have that

$$y_0 \notin] - \tilde{D}, \tilde{D}[\Rightarrow y(t; 0, x_0, y_0) \notin] - \tilde{d}, \tilde{d}[,$$

and

$$y_0 \in [-\tilde{D}, \tilde{D}] \Rightarrow y(t; 0, x_0, y_0) \in [-\tilde{\beta}, \tilde{\beta}],$$

for every $t \in [0, T]$ and $x_0 \in [0, 2\pi]$. Following the lines of the proof of Theorem 1, we can thus conclude that, if

$$2(\|E\|_\infty + \|h\|_1) < \gamma,$$

then equation (25) has at least two geometrically distinct T -periodic solutions. Moreover, in the spirit of Theorem 3, for every integer k such that

$$|k| < \frac{T}{2\pi} \phi^{-1}(\gamma - 2\|E\|_\infty - 2\|h\|_1),$$

equation (25) has at least two geometrically distinct periodic solutions of the second kind, satisfying

$$x(t + T) = x(t) + 2\pi k, \quad \text{for every } t \in \mathbb{R}.$$

Notice that, since adding a constant to $E(t)$ does not affect the original differential equation, we can always replace $\|E\|_\infty$ by $\frac{1}{2}(\max E - \min E)$.

As an example, we could take

$$\phi(s) = \frac{s}{\sqrt{1 + (s/\gamma)^2}}, \quad (26)$$

(here $c = +\infty$), leading to the mean curvature operator. We then get the following immediate consequence.

Corollary 2 *Assume that $e(t)$ satisfies (2) and let $a > 0$ and $\gamma > 0$ be such that $2(\|E\|_\infty + aT) < \gamma$. Then, the equation*

$$\frac{d}{dt} \frac{\dot{x}}{\sqrt{1 + (\dot{x}/\gamma)^2}} + a \sin x = e(t)$$

has at least two geometrically distinct T -periodic solutions. Moreover, for every integer k such that

$$|k| < \frac{T}{2\pi} \phi^{-1}(\gamma - 2\|E\|_\infty - 2aT),$$

there are at least two geometrically distinct periodic solutions of the second kind, satisfying

$$x(t + T) = x(t) + 2\pi k, \quad \text{for every } t \in \mathbb{R}.$$

The problem of subharmonic solutions of the second kind can be treated similarly, in the spirit of Theorem 4. For brevity, we prefer not entering into details.

This type of equations, with the mean curvature operator, has been studied using a variational method by Obersnel and Omari [29], who proved that, under the weaker condition $\|E\|_\infty < \gamma$, there are at least two geometrically distinct T -periodic “bounded variation” solutions. Here, however, we are always dealing with regular solutions. See also [35], where a different topological approach has been used. Notice that, as proved [29, Proposition 1.1], the zero mean condition (2) is not sufficient to ensure the existence of T -periodic solutions, so that some additional assumption is indeed needed.

References

- [1] V. I. Arnold, Problems in present day mathematics: fixed points of symplectic diffeomorphisms. Proc. Symp. Pure Math., vol. 28, Amer. Math. Soc., Providence 1976, p.66.
- [2] C. Bereanu and P. J. Torres, Existence of at least two periodic solutions of the forced relativistic pendulum. Proc. Amer. Math. Soc., to appear.
- [3] H. Brezis and J. Mawhin, Periodic solutions of the forced relativistic pendulum. Differential Integral Equations 23 (2010), 801–810.
- [4] H. Brezis and J. Mawhin, Periodic solutions of Lagrangian systems of relativistic oscillators. Communic. Applied Anal., to appear.
- [5] K. C. Chang, On the periodic nonlinearity and the multiplicity of solutions. Nonlinear Anal. 13 (1989), 527–537.

- [6] C. C. Conley and E. Zehnder, The Birkhoff–Lewis fixed point theorem and a conjecture of V. I. Arnold. *Invent. Math.* 73 (1983), 33–49.
- [7] C. C. Conley and E. Zehnder, Morse type index theory for flows and periodic solutions for Hamiltonian equations. *Comm. Pure Appl. Math.* 37 (1984), 207–253.
- [8] E. N. Dancer, On the use of asymptotics in nonlinear boundary value problems. *Ann. Mat. Pura Appl.* 131 (1982), 167–185.
- [9] W.-Y. Ding, A generalization of the Poincaré–Birkhoff theorem. *Proc. Amer. Math. Soc.* 88 (1983), 341–346.
- [10] A. Fonda and J. Mawhin, Multiple periodic solutions of conservative systems with periodic nonlinearity. In: *Differential equations and applications*, (Columbus, 1988), 298–304, Ohio Univ. Press, Athens, 1989.
- [11] A. Fonda and J. Mawhin, Critical point theory and multiple periodic solutions of conservative systems with periodic nonlinearity. In: *The Problem of Plateau*, 111–128, World Sci. Publ., River Edge, 1992.
- [12] A. Fonda, M. Sabatini and F. Zanolin, Periodic solutions of perturbed Hamiltonian systems in the plane by the use of the Poincaré–Birkhoff Theorem. *Topol. Meth. Nonlin. Anal.*, to appear.
- [13] A. Fonda and R. Toader, Periodic solutions of radially symmetric perturbations of Newtonian systems. *Proc. Amer. Math. Soc.*, to appear.
- [14] J. Franks, Generalizations of the Poincaré–Birkhoff theorem. *Ann. Math.* 128 (1988), 139–151.
- [15] J. Franks, Erratum to: "Generalizations of the Poincaré–Birkhoff theorem". *Ann. Math.* 164 (2006), 1097–1098.
- [16] F. Gantmacher, *Lectures in analytical mechanics*. MIR Publishers, Moscow, 1975.
- [17] G. Hamel, Ueber erzwungene Schwingungen bei endlichen Amplituden. *Math. Ann.* 86 (1922), 1–13.
- [18] N. Hingston, Subharmonic solutions of Hamiltonian equations on tori. *Ann. Math. (2)* 170 (2009), 529–560.
- [19] M. A. Krasnosel'skiĭ, *Translations Along Trajectories of Differential Equations*. American Mathematical Society, Providence, Rhode Island, 1968.
- [20] M. Levi, KAM theory for particles in periodic potentials. *Ergodic Theory Dynam. Systems* 10 (1990), 777–785.

- [21] R. Manasevich and J. Ward, On a result of Brezis and Mawhin. Proc. Amer. Math. Soc., to appear.
- [22] J. Mawhin, Forced second order conservative systems with periodic nonlinearity. *Analyse non linéaire* (Perpignan, 1987). Ann. Inst. H. Poincaré Anal. Non Linéaire 6 (1989), suppl., 415–434.
- [23] J. Mawhin, Nonlinear oscillations: one hundred years after Lyapunov and Poincaré. *Z. Angew. Math. Mech.* 73 (1993), T54–T62.
- [24] J. Mawhin, Global results for the forced pendulum equations. In: *Handbook of Differential Equations. Ordinary Differential Equations. Vol. I*, Elsevier, Amsterdam, 2004, pp. 533–589.
- [25] J. Mawhin, Multiplicity of solutions of variational systems involving ϕ -Laplacians with singular ϕ and periodic nonlinearities. *Discrete Contin. Dynam. Syst.*, to appear.
- [26] J. Mawhin and M. Willem, Multiple solutions of the periodic boundary value problem for some forced pendulum-type equations. *J. Differential Equations* 52 (1984), 264–287.
- [27] J. Mawhin and M. Willem, Variational methods and boundary value problems for vector second order differential equations and applications to the pendulum equation. In: *Nonlinear Analysis and Optimization (Bologna, 1982)*, 181–192, Lecture Notes in Math., 1107, Springer, Berlin, 1984.
- [28] J. Mawhin and M. Willem, *Critical Point Theory and Hamiltonian Systems*. Applied Mathematical Sciences, 74, Springer, Berlin, 1989.
- [29] F. Obersnel and P. Omari, Multiple bounded variation solutions of a periodically perturbed sine-curvature equation. *Commun. Contemp. Math.* 13 (2011), 1–21.
- [30] P. H. Rabinowitz, On a class of functionals invariant under a \mathbb{Z}^n action. *Trans. Amer. Math. Soc.* 310 (1988), 303–311.
- [31] C. Rebelo, A note on the Poincaré–Birkhoff fixed point theorem and periodic solutions of planar systems. *Nonlinear Anal.* 29 (1997), 291–311.
- [32] A. Szulkin, A relative category and applications to critical point theory for strongly indefinite functionals. *Nonlinear Anal.* 15 (1990), 725–739.
- [33] A. Szulkin, Cohomology and Morse theory for strongly indefinite functionals. *Math. Z.* 209 (1992), 375–418.
- [34] C.-L. Tang and X.-P. Wu, A note on periodic solutions of nonautonomous second-order systems. *Proc. Amer. Math. Soc.* 132 (2004), 1295–1303.

- [35] P. Torres, Nondegeneracy of the periodically forced Liénard differential equation with ϕ -Laplacian. *Comm. Contemporary Math.* 13 (2011), 283–292.
- [36] M. Willem, Oscillations forcées de l'équation du pendule. *Publ. IRMA Lille 3* (1981), v1–v3.
- [37] J. You, Invariant tori and Lagrange stability of pendulum-type equations. *J. Differential Equations* 85 (1990), 54–65.

Authors' addresses:

Alessandro Fonda
Dipartimento di Matematica e Informatica
Università di Trieste
P.le Europa 1
I-34127 Trieste
Italy
e-mail: a.fonda@units.it

Rodica Toader
Dipartimento di Ingegneria Civile e Architettura
Università di Udine
Via delle Scienze 206
I-33100 Udine
Italy
e-mail: toader@uniud.it

Mathematics Subject Classification: 34C25

Keywords: periodic solutions; pendulum equation; Poincaré–Birkhoff theorem; nonlinear dynamics.