

Nonlinear perturbations of some non-invertible differential operators

Alessandro Fonda and Rodica Toader

ABSTRACT. We consider perturbations, depending on a small parameter λ , of a non-invertible differential operator having a nonnegative spectrum. Given a pair of lower and upper solutions, belonging to the kernel of the differential operator, without any prescribed order, we prove the existence of a solution, when λ is sufficiently small. Our method of proof has the advantage to permit a uniform choice of λ for a whole class of functions. Applications are given in a variety of situations, ranging from ODE problems to equations of parabolic type, or involving the p -Laplacian operator.

1 Introduction

In this paper we deal with problems of the type

$$Lu = Nu,$$

where L is a non-invertible differential operator, and N is, in some sense, a nonlinear perturbation. A typical situation is that of a linear operator L , with compact resolvent, having zero as its first eigenvalue, and of a nonlinear operator

$$N = N_\lambda,$$

depending continuously on a small parameter λ , with $N_0 = 0$. Problems of this type are sometimes referred to as “nonlinear eigenvalue problems” (see, e.g., [3]). More generally, we can also consider some type of nonlinear differential operators.

To illustrate our results in a simple context, let us consider the following Neumann problem associated to an elliptic equation

$$\begin{cases} -\Delta u = \lambda f(x, u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where Ω is a bounded domain in \mathbb{R}^N with regular boundary, $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is L^r -Carathéodory, for some $r > N$, and λ is a real parameter.

Let us state the following theorem, which will be useful in order to compare our method with those used in the previous literature.

Theorem 1 *Assume that there exist two constants α, β such that*

$$f(x, \beta) \leq 0 \leq f(x, \alpha), \quad \text{a.e. in } \Omega.$$

Then, there is a $\bar{\lambda} > 0$ such that, for every $\lambda \in [0, \bar{\lambda}]$, problem (1) has a solution.

Notice that, in the above statement, α is a constant lower solution and β is an upper solution for (1). It is a classical result that a solution exists when α is a lower solution, β is an upper solution, and $\alpha \leq \beta$. When α and β are not ordered in this way, some interesting results have been obtained, for λ not necessarily small, assuming a growth condition on the nonlinearity which prevents the interaction with the positive part of the spectrum. See, e.g., [4, 29, 21, 22, 17, 18], for different kinds of equations.

Since λ is assumed to be small, Theorem 1 can be interpreted as a bifurcation type of result. In this framework, some general abstract theorems based on degree theory are already available, see e.g. [20, Theorem IV.2] or [19]. Roughly speaking, the main idea there is that, if the degree of some projected equation on the kernel is nonzero, it has to be nonzero also for a small perturbation.

We propose here a different type of approach, based on lower and upper solutions, which has the advantage to show that the choice of λ can be made uniformly for a whole class of functions f . For instance, we are able to prove the following generalization of Theorem 1.

Theorem 2 *Let $I \subseteq \mathbb{R}$ be an interval, and Λ be a positive number. Given a compact interval J , contained in the interior of I , there is a $\bar{\lambda} > 0$ with the following property: for every $\lambda \in [0, \bar{\lambda}]$ and every function f satisfying*

$$|f(x, u)| \leq h(x) \quad \text{for a.e. } x \in \Omega \text{ and every } u \in I,$$

with $\|h\|_{L^r} \leq \Lambda$, if there are two constants α, β in J for which

$$f(x, \beta) \leq 0 \leq f(x, \alpha) \quad \text{a.e. in } \Omega,$$

then problem (1) has a solution u , with $u(x) \in I$ for every $x \in \bar{\Omega}$.

The fact that $\bar{\lambda}$ can be chosen in such a uniform way is not evident from the above mentioned degree arguments, since it is not guaranteed that the associated degree remains nonzero for a small perturbation of the whole class of functions f considered in the statement.

Let us give a sketch of our method of proof. Let $\beta < \alpha$ be the constant upper and lower solutions. Modifying the function f outside the interval $[\beta - \varepsilon, \alpha + \varepsilon]$, for a small $\varepsilon > 0$, we can get a further pair of constant lower and upper solutions for the new problem, say $\beta - \delta$ and $\alpha + \delta$, for some $\delta > \varepsilon$. More precisely, $\beta - \delta$ is a strict lower solution, and $\alpha + \delta$ is a strict upper solution. So, degree theory applies, saying that the pair $(\beta - \delta, \alpha + \delta)$ is admissible and the associated degree's value is 1.

We now need to prove the following: *There is a solution between $\beta - \delta$ and $\alpha + \delta$, which enters the interval $[\beta, \alpha]$ somewhere.*

- If both the pairs $(\beta - \delta, \beta)$ and $(\alpha, \alpha + \delta)$ are admissible for the degree, then both the associated degree's values are 1 and, by the excision property, there must be a solution between $\beta - \delta$ and $\alpha + \delta$, which is neither between $\beta - \delta$ and β , nor between α and $\alpha + \delta$.

- Otherwise, assume for instance that the pair $(\beta - \delta, \beta)$ is not admissible for the degree. Being $\beta - \delta$ a strict lower solution, there must be a solution between $\beta - \delta$ and β which touches β somewhere. Similarly, if the pair $(\alpha, \alpha + \delta)$ is not admissible, there must be a solution between α and $\alpha + \delta$ which touches α somewhere.

At this point, having a solution between $\beta - \delta$ and $\alpha + \delta$, which enters the interval $[\beta, \alpha]$ somewhere, we use some classical estimates and the fact that λ is small to show that this solution cannot go far from the interval $[\beta, \alpha]$ and is indeed contained in $[\beta - \varepsilon, \alpha + \varepsilon]$, thus being a solution of the original problem.

Our ideas can be used in a large variety of situations, which do not need a variational structure. For instance, we can deal with the Neumann-periodic problem associated to a parabolic equation, or with the periodic problem for a second order ODE including friction terms. Also, some nonlinear differential operators can be considered, as for instance the p -Laplacian operator. Moreover, we will show how to deal with a wider class of nonlinearities, not necessarily involving an explicit dependence on a parameter λ .

We will consider the following three examples, which can be deduced from the above arguments, after a suitable rescaling. Here, $p > 1$ can be any exponent greater than 1.

I. The problem

$$\begin{cases} -\Delta u = |u|^{p-1}u + e(x) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases}$$

has a solution, provided that $\|e\|_{L^\infty}$ is sufficiently small.

II. The problem

$$\begin{cases} -\Delta u = (u^+)^p + e(x) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases}$$

has a solution, provided that $e(x) \leq 0$ for a.e. $x \in \Omega$, and $\|e\|_{L^\infty}$ is small enough. Here, $u^+ = \max\{u, 0\}$.

III. The problem

$$\begin{cases} -\Delta u = u^p - \lambda a(x)u^q & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases}$$

has a solution, provided that $a(x)$ is continuous and positive, $q < p$, and $\lambda > 0$ is sufficiently small.

Our method, moreover, permits to relate the smallness of the forcing term to the powers p, q . For instance, in the above examples I and II, if p is sufficiently large, any function e such that

$$\|e\|_{L^\infty} \leq \frac{1}{2^p}$$

will guarantee the existence of a solution.

Similar problems, with various kinds of boundary conditions, have been studied by many authors, mainly by the use of variational methods. See, e.g., [33, 6, 7, 32, 36, 11, 8, 5, 9, 24] for the Dirichlet boundary conditions and [23, 1, 35, 13, 12] for the Neumann problem. The main difficulty, in this framework, is the loss of compactness encountered at the critical exponent $p = 2^* - 1 = \frac{N+2}{N-2}$, when $N \geq 3$.

The paper is organized as follows. In Section 2 we consider the Neumann-periodic problem associated to a parabolic equation

$$\begin{cases} \partial_t u - \Delta u = \lambda f(x, t, u, \nabla u) & \text{in } \Omega \times]0, T[, \\ \partial_\nu u = 0 & \text{on } \partial\Omega \times]0, T[, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases}$$

assuming a Bernstein-Nagumo type of condition in order to deal with the dependence on ∇u . We provide in full details the proof of our result for this case, which presents some more technical aspects, so that in the rest of the paper we will only need to show how to modify our arguments in order to deal with different kinds of problems.

A linear elliptic operator is considered in Section 3. The same arguments used for the parabolic equation are easily adapted to this case.

In Section 4, a second order ODE is studied, where more specific conditions of Bernstein-Nagumo type can be considered. Equations of Liénard or Rayleigh type are also treated there. We thus generalize a result stated in [31].

In Section 5 we deal with the p -Laplacian operator. Since a weaker notion of solution is adopted, a more detailed exposition of the proof is given in this case.

For the reader's convenience, we provide in the Appendix a proof of the fundamental estimate used, in connection with the Bernstein-Nagumo condition, in the ODE case treated in Section 4.

2 The parabolic equation

Let Ω be a bounded domain in \mathbb{R}^N with a C^2 -boundary $\partial\Omega$. Given $T > 0$, set $Q_T = \Omega \times]0, T[$ and $\Sigma_T = \partial\Omega \times]0, T[$.

Define the elliptic differential operator

$$\mathcal{A}u := - \sum_{i,j=1}^N a_{ij}(x, t) \partial_{x_i x_j}^2 u + \sum_{i=1}^N a_i(x, t) \partial_{x_i} u.$$

Here $a_{ij} \in C(\overline{Q_T})$, $a_{ij} = a_{ji}$, $a_{ij}(x, 0) = a_{ij}(x, T)$ in $\overline{\Omega}$, for $i, j = 1, \dots, N$, there exists $\bar{a} > 0$ such that

$$\sum_{i,j=1}^N a_{ij}(x, t) \xi_i \xi_j \geq \bar{a} |\xi|^2, \quad \text{for every } (x, t, \xi) \in \overline{Q_T} \times \mathbb{R}^N,$$

and $a_i \in L^\infty(Q_T)$, for $i = 1, \dots, N$.

We consider the following Neumann-periodic problem

$$(P) \quad \begin{cases} \partial_t u + \mathcal{A}u = f(x, t, u, \nabla_x u) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega. \end{cases}$$

Here f is a L^r -Carathéodory function, for some $r > N + 2$, ν is the unit outer normal to $\partial\Omega$ and $\partial_\nu u = \sum_{i=1}^N \partial_{x_i} u \nu_i$ is the normal derivative of u . If necessary, all functions defined on Q_T will be assumed to be extended by T -periodicity to $\Omega \times \mathbb{R}$.

Let $W_r^{2,1}(Q_T)$ be the space of functions u such that

$$u, \partial_t u, \partial_{x_i} u, \partial_{x_i x_j}^2 u \in L^r(Q_T),$$

for $i, j = 1, \dots, N$, with the usual norm

$$\|u\|_{W_r^{2,1}} = \|u\|_{L^r} + \|\partial_t u\|_{L^r} + \sum_{i=1}^N \|\partial_{x_i} u\|_{L^r} + \sum_{i,j=1}^N \|\partial_{x_i x_j}^2 u\|_{L^r}.$$

Recall that $W_r^{2,1}(Q_T)$ is compactly imbedded into $C^{1,0}(\overline{Q_T})$.

We say that u is a *solution* of problem (P) if u belongs to $W_r^{2,1}(Q_T)$, it satisfies the differential equation a.e. in Q_T and the boundary and periodicity conditions pointwise. A function with these properties is usually called “strong solution” in the literature.

Let us define the class of functions

$$\mathcal{F}(I, \Lambda, K),$$

where $I \subseteq \mathbb{R}$ is an interval, and Λ, K are some nonnegative constants. Its elements are the Carathéodory functions $f : Q_T \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ which satisfy the following Bernstein-Nagumo growth condition:

$$(BN) \quad \begin{aligned} &|f(x, t, u, v)| \leq h(x, t) + K|v|^2, \\ &\text{for a.e. } (x, t) \in Q_T \text{ and every } (u, v) \in I \times \mathbb{R}^N, \end{aligned}$$

for some $h \in L^r(Q_T)$, with $r > N + 2$ and $\|h\|_{L^r} \leq \Lambda$.

The following classical result, whose proof for our setting can be found in [18, Proposition III.1.4.], will be used in connection with the Bernstein-Nagumo condition, in order to get uniform estimates on the norm of the solutions.

Proposition 1 (Fundamental estimate) *Given $M, \Lambda, K > 0$, there is a constant $C > 0$ such that, if $u \in W_r^{2,1}(Q_T)$ verifies*

$$\begin{cases} |\partial_t u + \mathcal{A}u| \leq h(x, t) + K|\nabla_x u|^2, & \text{a.e. in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ |u| \leq M & \text{in } Q_T, \end{cases}$$

for some $h \in L^r(Q_T)$ with $\|h\|_{L^r} \leq \Lambda$, then

$$\|u\|_{W_r^{2,1}} \leq C.$$

Let us now state our main result.

Theorem 3 *Let $I \subseteq \mathbb{R}$ be an interval. Given a compact interval J , contained in the interior of I , there is a $\Lambda > 0$ such that, for every $f \in \mathcal{F}(I, \Lambda, \Lambda)$, if there are two constants α, β in J for which*

$$f(x, t, \beta, 0) \leq 0 \leq f(x, t, \alpha, 0) \quad \text{a.e. in } Q_T,$$

then problem (P) has a solution u , with $u(x, t) \in I$ for every $(x, t) \in \overline{Q}_T$.

In order to prove the theorem we need some preliminary considerations.

Definition 1 A function $\alpha \in W_r^{2,1}(Q_T)$ is a lower solution of (P) if

$$\begin{cases} \partial_t \alpha + \mathcal{A}\alpha \leq f(x, t, \alpha, \nabla_x \alpha) & \text{a.e. in } Q_T, \\ \partial_\nu \alpha \leq 0 & \text{on } \Sigma_T, \\ \alpha(x, 0) \leq \alpha(x, T) & \text{in } \Omega. \end{cases}$$

The function α is a strict lower solution if it is a lower solution and, for every solution u of (P) with $u \geq \alpha$, one has that $u > \alpha$ on \overline{Q}_T .

Analogously, a function $\beta \in W_r^{2,1}(Q_T)$ is an upper solution of (P) if

$$\begin{cases} \partial_t \beta + \mathcal{A}\beta \geq f(x, t, \beta, \nabla_x \beta) & \text{a.e. in } Q_T, \\ \partial_\nu \beta \geq 0 & \text{on } \Sigma_T, \\ \beta(x, 0) \geq \beta(x, T) & \text{in } \Omega. \end{cases}$$

The function β is a strict upper solution if it is an upper solution and, for every solution u of (P) with $u \leq \beta$, one has that $u < \beta$ on \overline{Q}_T .

Notice that the constants α and β appearing in the statement of Theorem 3 are lower and upper solutions, respectively, although they might not be strict.

Let us introduce the linear operator

$$L : D(L) \subset W_r^{2,1}(Q_T) \rightarrow L^r(Q_T),$$

where

$$D(L) = \{u \in W_r^{2,1}(Q_T) : \partial_\nu u = 0 \text{ on } \Sigma_T \text{ and } u(x, 0) = u(x, T) \text{ in } \Omega\},$$

defined by

$$Lu = \partial_t u + \mathcal{A}u.$$

The operator L is not invertible: 0 is its first eigenvalue, and $\ker L$ is made of the constant functions. Let us fix a real number $\sigma > 0$. One can prove (cf. [26, Lemma 4.1 and Corollary 5.6], see also [18, Proposition I.1.3]) that $L + \sigma I$ is invertible, with a continuous inverse

$$(L + \sigma I)^{-1} : L^r(Q_T) \rightarrow W_r^{2,1}(Q_T).$$

Let us also introduce the continuous nonlinear operator

$$N : C^{1,0}(\overline{Q}_T) \rightarrow L^r(Q_T),$$

defined by

$$(Nu)(x, t) = f(x, t, u(x, t), \nabla_x u(x, t)).$$

Problem (P) can be written as

$$Lu = Nu,$$

which is equivalent to the fixed point problem

$$u = \mathcal{S}u,$$

where the function

$$\mathcal{S} : C^{1,0}(\overline{Q}_T) \rightarrow C^{1,0}(\overline{Q}_T)$$

is defined by

$$\mathcal{S}u = (L + \sigma I)^{-1}(N + \sigma I)u.$$

Since $W_r^{2,1}(Q_T)$ is compactly imbedded in $C^{1,0}(\overline{Q}_T)$, we have that \mathcal{S} is completely continuous, so that we can use Leray-Schauder degree theory.

Definition 2 *Let $\alpha, \beta : \overline{Q}_T \rightarrow \mathbb{R}$ be two continuous functions. The pair (α, β) is said to be admissible for (P) if $\alpha < \beta$ and there is $R > 0$ with the following property: any solution u of (P) satisfying $\alpha \leq u \leq \beta$ is such that*

$$\alpha < u < \beta \quad \text{and} \quad \|u\|_{C^{1,0}} < R.$$

Notice that, if (α, β) is an admissible pair, then the set

$$\mathcal{U}_{(\alpha, \beta)} := \{u \in C^{1,0}(\overline{Q}_T) : \alpha < u < \beta\}$$

is open, nonempty, and any fixed point of \mathcal{S} contained in $\mathcal{U}_{(\alpha, \beta)}$ belongs to the ball B_R in $C^{1,0}(\overline{Q}_T)$. Hence, we can define

$$\deg(I - \mathcal{S}, \mathcal{U}_{(\alpha, \beta)}) = d_{\text{LS}}(I - \mathcal{S}, \mathcal{U}_{(\alpha, \beta)} \cap B_R),$$

where d_{LS} denotes the Leray-Schauder degree.

The following classical result on lower and upper solutions has been proved in this setting in [18, Theorem II.2.1], although we state it here in a slightly more general form.

Proposition 2 (Well-ordered lower and upper solutions) *In the above setting, assume that α is a lower solution and β is an upper solution of (P) satisfying $\alpha \leq \beta$. Then problem (P) has a solution u such that $\alpha \leq u \leq \beta$. If, moreover, (α, β) is an admissible pair, then*

$$\deg(I - \mathcal{S}, \mathcal{U}_{(\alpha, \beta)}) = 1.$$

Proof of Theorem 3. Let $\delta > 0$ be such that $[\min J - \delta, \max J + \delta] \subseteq I$, and fix $\varepsilon \in]0, \frac{\delta}{2}[$. We will prove the statement by taking $\Lambda = \frac{1}{m}$, with $m \geq 1$ a sufficiently large integer. By contradiction, assume there exist a function $f_m \in \mathcal{F}(I, \frac{1}{m}, \frac{1}{m})$, and two constants $\alpha_m, \beta_m \in J$, satisfying

$$f_m(x, t, \beta_m, 0) \leq 0 \leq f_m(x, t, \alpha_m, 0) \quad \text{a.e. in } Q_T,$$

for which the problem

$$(P_m) \quad \begin{cases} \partial_t u + \mathcal{A}u = f_m(x, t, u, \nabla_x u) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases}$$

has no solution u , with $u(x, t) \in I$ for every $(x, t) \in \overline{Q}_T$. By Proposition 2, it has to be $\beta_m < \alpha_m$. Define the modified function

$$\tilde{f}_m(x, t, u, v) = \begin{cases} \frac{1}{m} & \text{if } u \leq \beta_m - 2\varepsilon, \\ \frac{1}{m} + \frac{1}{\varepsilon}(f_m(x, t, u, v) - \frac{1}{m})(u - \beta_m + 2\varepsilon), & \text{if } \beta_m - 2\varepsilon \leq u \leq \beta_m - \varepsilon, \\ f_m(x, t, u, v) & \text{if } \beta_m - \varepsilon \leq u \leq \alpha_m + \varepsilon, \\ -\frac{1}{m} + \frac{1}{\varepsilon}(f_m(x, t, u, v) + \frac{1}{m})(\alpha_m + 2\varepsilon - u), & \text{if } \alpha_m + \varepsilon \leq u \leq \alpha_m + 2\varepsilon, \\ -\frac{1}{m} & \text{if } u \geq \alpha_m + 2\varepsilon, \end{cases}$$

and consider the problem

$$(\tilde{P}_m) \quad \begin{cases} \partial_t u + \mathcal{A}u = \tilde{f}_m(x, t, u, \nabla_x u) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega. \end{cases}$$

Notice that $\tilde{f}_m \in \mathcal{F}(\mathbb{R}, \frac{1}{m}(1 + |Q_T|^{1/r}), \frac{1}{m})$. Let us prove that, for the modified problem (\tilde{P}_m) ,

$$\beta_m - \delta \text{ is a strict lower solution.}$$

Indeed, as $\tilde{f}_m(x, t, \beta_m - \delta, 0) = \frac{1}{m}$, clearly $\beta_m - \delta$ is a lower solution. If it were not strict, there would be a solution u of (\tilde{P}_m) with $u \geq \beta_m - \delta$, and a point $(x_0, t_0) \in \overline{Q}_T$ for which $u(x_0, t_0) = \beta_m - \delta$. Setting $v = u - \beta_m + \delta$, there is $\rho > 0$ such that

$$\partial_t v + \mathcal{A}v = \frac{1}{m} > 0 \quad \text{a.e. in } (B(x_0, \rho) \cap \Omega) \times]t_0 - \rho, t_0 + \rho[.$$

If $x_0 \in \Omega$, we take $\rho > 0$ small enough so that $B(x_0, \rho)$ is contained in Ω and, by the Strong Maximum Principle (see, e.g., [18, Proposition I.1.1]), it has to be $v = 0$ in $B(x_0, \rho) \times]t_0 - \rho, t_0]$, which is clearly impossible. On the other hand, if $x_0 \in \partial\Omega$, it should be $\partial_\nu v(x_0, t_0) < 0$, in contradiction with the Neumann boundary condition.

In the same way we can show that

$\alpha_m + \delta$ is a strict upper solution.

Hence, the pair $(\beta_m - \delta, \alpha_m + \delta)$ is admissible for (\tilde{P}_m) . Indeed, let u be a solution of (\tilde{P}_m) with $\beta_m - \delta \leq u \leq \alpha_m + \delta$. Since $\tilde{f}_m \in \mathcal{F}(\mathbb{R}, 1 + |Q_T|^{1/r}, 1)$, by Proposition 1 there is a constant $R > 0$ such that $\|u\|_{C^{1,0}} < R$. Since $\beta_m - \delta$ and $\alpha_m + \delta$ are strict, we also have that $\beta_m - \delta < u < \alpha_m + \delta$.

Let us introduce the operators $\tilde{N}_m : C^{1,0}(\overline{Q}_T) \rightarrow L^r(Q_T)$, defined by

$$(\tilde{N}_m u)(x, t) = \tilde{f}_m(x, t, u(x, t), \nabla_x u(x, t)),$$

and $\tilde{\mathcal{S}}_m : C^{1,0}(\overline{Q}_T) \rightarrow C^{1,0}(\overline{Q}_T)$, defined by

$$\tilde{\mathcal{S}}_m u = (L + \sigma I)^{-1}(\tilde{N}_m + \sigma I)u. \quad (2)$$

Here, σ is any fixed positive number. We now distinguish two cases.

Case 1. The pairs $(\beta_m - \delta, \beta_m)$ and $(\alpha_m, \alpha_m + \delta)$ are admissible for (\tilde{P}_m) . By Proposition 2, we have

$$\begin{aligned} \deg(I - \tilde{\mathcal{S}}_m, \mathcal{U}_{(\beta_m - \delta, \beta_m)}) &= 1, \\ \deg(I - \tilde{\mathcal{S}}_m, \mathcal{U}_{(\alpha_m, \alpha_m + \delta)}) &= 1, \\ \deg(I - \tilde{\mathcal{S}}_m, \mathcal{U}_{(\beta_m - \delta, \alpha_m + \delta)}) &= 1. \end{aligned}$$

By the excision property of the degree,

$$\deg(I - \tilde{\mathcal{S}}_m, \mathcal{U}_{(\beta_m - \delta, \alpha_m + \delta)} \setminus \overline{\mathcal{U}_{(\beta_m - \delta, \beta_m)} \cup \mathcal{U}_{(\alpha_m, \alpha_m + \delta)}}) = -1.$$

Hence, there is a solution u_m of (\tilde{P}_m) such that

$$\beta_m - \delta < u_m < \alpha_m + \delta, \quad (3)$$

and, for some $(x_m, t_m) \in \overline{Q}_T$,

$$\beta_m \leq u_m(x_m, t_m) \leq \alpha_m. \quad (4)$$

Case 2. One or both of the pairs $(\beta_m - \delta, \beta_m)$ and $(\alpha_m, \alpha_m + \delta)$ are not admissible for (\tilde{P}_m) . Assume for instance $(\alpha_m, \alpha_m + \delta)$ is not admissible. Then, for every $R > 0$ there is a solution u_m of (\tilde{P}_m) with $\alpha_m \leq u_m \leq \alpha_m + \delta$ such that, either $\min(u_m - \alpha_m) = 0$, or $\min(\alpha_m + \delta - u_m) = 0$, or $\|u_m\|_{C^{1,0}} \geq R$. Since $\alpha_m + \delta$ is strict, we have $u_m < \alpha_m + \delta$. Since $\tilde{f}_m \in \mathcal{F}(\mathbb{R}, 1 + |Q_T|^{1/r}, 1)$, by Proposition 1 the third possibility cannot occur for R large enough, so that it has to be $\min(u_m - \alpha_m) = 0$. If instead $(\beta_m - \delta, \beta_m)$ is not admissible, analogously we find a solution u_m such that $\max(\beta_m - u_m) = 0$.

In both cases, we thus find a solution u_m of (\tilde{P}_m) verifying (3) and (4).

Since $\tilde{f}_m \in \mathcal{F}(\mathbb{R}, 1 + |Q_T|^{1/r}, 1)$, by Proposition 1 there is a constant $C > 0$ such that

$$\|u_m\|_{W_r^{2,1}} \leq C,$$

for every m . Being $W_r^{2,1}(Q_T)$ compactly imbedded in $C^{1,0}(\overline{Q}_T)$, there are a subsequence, still denoted by $(u_m)_m$, and a function $\bar{u} \in C^{1,0}(\overline{Q}_T)$ such that $u_m \rightarrow \bar{u}$ in $C^{1,0}(\overline{Q}_T)$. In particular, for every $(x, t) \in \overline{Q}_T$ we have $|\nabla_x u_m(x, t)| \leq \bar{c}$, for some constant $\bar{c} > 0$. As $\tilde{f}_m \in \mathcal{F}(\mathbb{R}, \frac{1}{m}(1 + |Q_T|^{1/r}), \frac{1}{m})$, by the Lebesgue dominated convergence Theorem we have

$$\tilde{N}_m u_m = \tilde{f}_m(\cdot, u_m(\cdot), \nabla_x u_m(\cdot)) \rightarrow 0 \quad \text{in } L^r(Q_T).$$

Then,

$$\tilde{\mathcal{S}}_m u_m = (L + \sigma I)^{-1}(\tilde{N}_m u_m + \sigma u_m) \rightarrow (L + \sigma I)^{-1}(\sigma \bar{u}) \quad \text{in } C^{1,0}(\overline{Q}_T).$$

As $u_m = \tilde{\mathcal{S}}_m u_m$, we get

$$\bar{u} = (L + \sigma I)^{-1}(\sigma \bar{u}),$$

i.e., $\bar{u} \in D(L)$ and $L\bar{u} = 0$. Hence, \bar{u} is constant. Since $u_m \rightarrow \bar{u}$ uniformly in \overline{Q}_T , for m large enough,

$$\text{osc}(u_m) = \max u_m - \min u_m \leq \varepsilon.$$

Recalling that u_m satisfies (4), it has to be

$$\beta_m - \varepsilon \leq u_m \leq \alpha_m + \varepsilon.$$

So, u_m is a solution to problem (P_m) and, since α_m, β_m are in J , by the choice of ε we have that $u_m(x, t) \in I$, for every $(x, t) \in \overline{Q}_T$. We thus get a contradiction, which ends the proof. \blacksquare

Remark 1 For Λ small enough, the solution provided by Theorem 3 has a small oscillation. This can be proved by contradiction, as follows. Assume that there exist $\bar{\varepsilon} > 0$, a sequence of functions $f_m \in \mathcal{F}(I, \frac{1}{m}, \frac{1}{m})$, satisfying the assumptions of the theorem, and a sequence of solutions u_m to problem (P_m) , with $u_m(x, t) \in I$ for every $(x, t) \in \overline{Q}_T$, for which $\text{osc}(u_m) > \bar{\varepsilon}$. Arguing as in the last part of the proof of Theorem 3, for a subsequence we have that $(u_m)_m$ converges to a constant \bar{u} , thus getting the contradiction.

Remark 2 Besides the Neumann condition one could consider more general oblique derivative boundary conditions, like

$$\sum_{i=1}^N b_i(x, t) \partial_{x_i} u = 0 \quad \text{on } \Sigma_T,$$

where $b_i \in C^{1,1/2}(\Sigma_T)$, $b_i(x, 0) = b_i(x, T)$ on $\partial\Omega$, for $i = 1, \dots, N$ and there is $\bar{b} > 0$ such that, for all $(x, t) \in \Sigma_T$,

$$\sum_{i=1}^N b_i(x, t) \nu_i(x) \geq \bar{b}.$$

Remark 3 Theorem 3 can be reformulated for the Dirichlet-periodic problem

$$(D) \quad \begin{cases} \partial_t u + \mathcal{A}u - \lambda_1 u = f(x, t, u, \nabla_x u) & \text{in } Q_T, \\ u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases}$$

where λ_1 denotes the first eigenvalue of the differential operator. The lower and upper solutions, in this case, will not be constant, but of the type

$$\alpha(x, t) = a \varphi_1(x, t), \quad \beta(x, t) = b \varphi_1(x, t),$$

where $\varphi_1(x, t)$ is the eigenfunction corresponding to λ_1 .

As a direct consequence of Theorem 3, we have the following.

Corollary 1 *Let $I \subseteq \mathbb{R}$ be an interval, and Λ, K be some fixed positive numbers. Given a compact interval J , contained in the interior of I , there is a $\bar{\lambda} > 0$ with the following property: for every $\lambda \in [0, \bar{\lambda}]$ and every $f \in \mathcal{F}(I, \Lambda, K)$, if there are two constants α, β in J for which*

$$f(x, t, \beta, 0) \leq 0 \leq f(x, t, \alpha, 0) \quad \text{a.e. in } Q_T,$$

then the problem

$$\begin{cases} \partial_t u + \mathcal{A}u = \lambda f(x, t, u, \nabla_x u) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases}$$

has a solution u , with $u(x, t) \in I$ for every $(x, t) \in \overline{Q}_T$.

As a particular case, we state the following.

Corollary 2 *Assume that $g : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function and $[a, b]$ is an interval such that*

$$[a, b] \subseteq g(\mathbb{R}).$$

Then, there is a $\bar{\lambda} > 0$ such that, for every $\lambda \in [0, \bar{\lambda}]$ and every $e \in L^\infty(Q_T)$ such that

$$e(x, t) \in [-b, -a], \quad \text{a.e. in } Q_T, \quad (5)$$

the problem

$$\begin{cases} \partial_t u + \mathcal{A}u = \lambda(g(u) + e(x, t)) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases} \quad (6)$$

has a solution.

Proof Let α and β be such that $g(\alpha) = b$ and $g(\beta) = a$. By (5), α is a lower solution and β is an upper solution of (6). If $\alpha \leq \beta$, the result is well-known. Assume $\beta < \alpha$ and take $I = [\beta - 1, \alpha + 1]$. Then, writing $f(x, t, u) = g(u) + e(x, t)$, we have that $f \in \mathcal{F}(I, \Lambda, 0)$, where

$$\Lambda = \max\{|g(s)| : s \in I\} + \max\{|a|, |b|\}.$$

Corollary 1 then gives the conclusion. \blacksquare

In particular, when the function g is unbounded from above and from below, any bounded function $e(x, t)$ will satisfy the assumptions.

Let us now provide two examples where our main result applies.

Corollary 3 *Let $\zeta \in L^\infty(Q_T)$ be a function bounded below by a positive constant. Given $p > 1$, the problem*

$$\begin{cases} \partial_t u + \mathcal{A}u = \zeta(x, t)|u|^{p-1}u + e(x, t) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases} \quad (7)$$

has a solution, provided that $\|e\|_{L^\infty}$ is sufficiently small. More precisely, for every given $\delta > 0$ there is a $\bar{\Lambda}_\delta > 0$ such that, for every $p > 1$, if

$$\|e\|_{L^\infty} \leq \left(\frac{\bar{\Lambda}_\delta}{(1 + \delta)^p} \right)^{\frac{p}{p-1}}, \quad (8)$$

then (7) has a solution.

Proof Fix $\delta > 0$ and consider the problem

$$\begin{cases} \partial_t w + \mathcal{A}w = \lambda(\zeta(x, t)|w|^{p-1}w + \tilde{e}(x, t)) & \text{in } Q_T, \\ \partial_\nu w = 0 & \text{on } \Sigma_T, \\ w(x, 0) = w(x, T) & \text{in } \Omega, \end{cases} \quad (9)$$

where λ is a positive constant and \tilde{e} is such that $\|\tilde{e}\|_{L^\infty} \leq \text{ess inf } \zeta$. Take $J = [-1, 1]$ and $I = [-1 - \delta, 1 + \delta]$. Let $\Lambda = \Lambda_\delta > 0$ be as in the statement of Theorem 3. Setting $f_\lambda(x, t, w) = \lambda(\zeta(x, t)|w|^{p-1}w + \tilde{e}(x, t))$, we have

$$f_\lambda(x, t, -1) \leq 0 \leq f_\lambda(x, t, 1),$$

for a.e. $(x, t) \in Q_T$ and, for every $w \in I$,

$$|f_\lambda(x, t, w)| \leq \lambda(\|\zeta\|_{L^\infty}(1 + \delta)^p + \text{ess inf } \zeta) \leq 2\lambda\|\zeta\|_{L^\infty}(1 + \delta)^p. \quad (10)$$

By Theorem 3, if

$$2\lambda\|\zeta\|_{L^\infty}(1 + \delta)^p |Q_T|^{1/r} \leq \Lambda_\delta, \quad (11)$$

then (9) has a solution w such that $w(x, t) \in I$, for every $(x, t) \in \overline{Q_T}$. Set

$$\bar{\Lambda}_\delta = \frac{\Lambda_\delta \min\{\text{ess inf } \zeta, 1\}}{2\|\zeta\|_{L^\infty} |Q_T|^{1/r}}.$$

Consider now (7), assume (8) with $\bar{\Lambda}_\delta > 0$ as above, and fix

$$\lambda = \frac{\Lambda_\delta}{2\|\zeta\|_{L^\infty}(1 + \delta)^p |Q_T|^{1/r}}, \quad (12)$$

so that (11) is satisfied. Setting $w = \lambda^{\frac{1}{1-p}} u$, we see that (7) is equivalent to (9), with $\tilde{e}(x, t) = \lambda^{\frac{p}{1-p}} e(x, t)$. Since, using (8) and (12),

$$\|\tilde{e}\|_{L^\infty} = \lambda^{\frac{p}{1-p}} \|e\|_{L^\infty} \leq \lambda^{\frac{p}{1-p}} \left(\frac{\bar{\Lambda}_\delta}{(1 + \delta)^p} \right)^{\frac{p}{p-1}} = \min\{\text{ess inf } \zeta, 1\}^{\frac{p}{p-1}} \leq \text{ess inf } \zeta,$$

we have that (9) is solvable. Therefore, (7) is solvable, as well. \blacksquare

Remark 4 Notice that, taking $\delta = \frac{1}{2}$, condition (8) is satisfied if p is sufficiently large and

$$\|e\|_{L^\infty} \leq \frac{1}{2^p}.$$

In a similar way, we have the following result, where we use the notation

$$u^+ = \max\{u, 0\}.$$

Corollary 4 *Let $\zeta \in L^\infty(Q_T)$ be a function bounded below by a positive constant. Given $p > 1$, the problem*

$$\begin{cases} \partial_t u + \mathcal{A}u = \zeta(x, t)(u^+)^p + e(x, t) & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega, \end{cases} \quad (13)$$

has a solution, provided that $e(x, t) \leq 0$ for almost every $(x, t) \in Q_T$, and $\|e\|_{L^\infty}$ is sufficiently small. More precisely, for every given $\delta > 0$ there is a $\bar{\Lambda}_\delta > 0$ such that, for every $p > 1$, if $e(x, t) \leq 0$ for almost every $(x, t) \in Q_T$ and (8) holds, then (13) has a solution.

Proof As in the proof of Corollary 3, first consider the problem

$$\begin{cases} \partial_t w + \mathcal{A}w = \lambda(\zeta(x, t)(w^+)^p + \tilde{e}(x, t)) & \text{in } Q_T, \\ \partial_\nu w = 0 & \text{on } \Sigma_T, \\ w(x, 0) = w(x, T) & \text{in } \Omega, \end{cases} \quad (14)$$

where \tilde{e} is such that $-\text{ess inf } \zeta \leq \tilde{e}(x, t) \leq 0$, for almost every $(x, t) \in Q_T$. Once $\delta > 0$ is fixed, take $J = [0, 1]$ and $I = [-\delta, 1 + \delta]$. Setting $f_\lambda(x, t, w) = \lambda(\zeta(x, t)(w^+)^p + \tilde{e}(x, t))$, we have

$$f_\lambda(x, t, 0) \leq 0 \leq f_\lambda(x, t, 1),$$

and (10) holds, for a.e. $(x, t) \in Q_T$ and every $w \in I$. One then concludes as in the proof of Corollary 3. \blacksquare

As a further example, let us consider the problem

$$\begin{cases} \partial_t u + \mathcal{A}u = \zeta_1(x, t)u^p - \lambda\zeta_2(x, t)u^q & \text{in } Q_T, \\ u > 0 & \text{in } Q_T, \\ \partial_\nu u = 0 & \text{on } \Sigma_T, \\ u(x, 0) = u(x, T) & \text{in } \Omega. \end{cases} \quad (15)$$

Corollary 5 *Assume that $\zeta_1, \zeta_2 \in L^\infty(Q_T)$ are bounded below by a positive constant, $p > 1$, and $q < p$. Then, there is a $\bar{\lambda} > 0$ such that, for every $\lambda \in [0, \bar{\lambda}]$, problem (15) has a solution.*

Proof Let us first show that, for $\hat{\lambda} > 0$ small enough, there is a positive solution of

$$\begin{cases} \partial_t w + \mathcal{A}w = \hat{\lambda}(\zeta_1(x, t)w^p - \zeta_2(x, t)w^q) & \text{in } Q_T, \\ w > 0 & \text{in } Q_T, \\ \partial_\nu w = 0 & \text{on } \Sigma_T, \\ w(x, 0) = w(x, T) & \text{in } \Omega. \end{cases} \quad (16)$$

Let α, β be defined by

$$\alpha = \left(\frac{\text{ess sup } \zeta_2}{\text{ess inf } \zeta_1} \right)^{\frac{1}{p-q}}, \quad \beta = \left(\frac{\text{ess inf } \zeta_2}{\text{ess sup } \zeta_1} \right)^{\frac{1}{p-q}}.$$

Then $0 < \beta \leq \alpha$, and

$$\zeta_1(x, t)\beta^p - \zeta_2(x, t)\beta^q \leq 0 \leq \zeta_1(x, t)\alpha^p - \zeta_2(x, t)\alpha^q,$$

for a.e. $(x, t) \in Q_T$. Taking $\varepsilon \in]0, \beta[$ and applying Corollary 1, we obtain a positive solution w of (16), for $\hat{\lambda} > 0$ sufficiently small.

It is now easy to see that the rescaled function $u = \hat{\lambda}^{\frac{1}{p-1}} w$ is a solution of (15), with $\lambda = \hat{\lambda}^{\frac{p-q}{p-1}}$. \blacksquare

3 The elliptic equation

In this section, we consider the Neumann problem

$$\begin{cases} -\sum_{i,j=1}^N a_{ij}(x) \partial_{x_i x_j}^2 u + \sum_{i=1}^N a_i(x) \partial_{x_i} u = f(x, u, \nabla u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (17)$$

Here, Ω is a bounded domain in \mathbb{R}^N with a C^2 -boundary, $a_{ij} \in C(\overline{\Omega})$, $a_{ij} = a_{ji}$, for $i, j = 1, \dots, N$, there exists $\bar{a} > 0$ such that

$$\sum_{i,j=1}^N a_{ij}(x) \xi_i \xi_j \geq \bar{a} |\xi|^2, \quad \text{for every } (x, \xi) \in \overline{\Omega} \times \mathbb{R}^N,$$

and $a_i \in L^\infty(\Omega)$, for $i = 1, \dots, N$. The function $f : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ is assumed to be L^r -Carathéodory, for some $r > N$.

We say that u is a *solution* of problem (17) if u belongs to $W^{2,r}(\Omega)$, it satisfies the differential equation a.e. in Ω and the boundary condition pointwise. (Here and in the sequel we use the standard notation for the Sobolev space $W^{2,r}(\Omega)$, which should not be confused with the notations used in the previous section.)

Let us define the class of functions

$$\mathcal{F}(I, \Lambda, K),$$

where $I \subseteq \mathbb{R}$ is an interval, and Λ, K are some nonnegative constants. Its elements are the Carathéodory functions $f : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ which satisfy the following Bernstein-Nagumo growth condition:

$$(BN) \quad \begin{aligned} |f(x, u, v)| &\leq h(x) + K|v|^2, \\ \text{for a.e. } x \in \Omega \text{ and every } (u, v) &\in I \times \mathbb{R}^N, \end{aligned}$$

for some $h \in L^r(\Omega)$, with $r > N$ and $\|h\|_{L^r} \leq \Lambda$.

Theorem 4 *Let $I \subseteq \mathbb{R}$ be an interval. Given a compact interval J , contained in the interior of I , there is a $\Lambda > 0$ such that, for every $f \in \mathcal{F}(I, \Lambda, \Lambda)$, if there are two constants α, β in J for which*

$$f(x, \beta, 0) \leq 0 \leq f(x, \alpha, 0) \quad \text{a.e. in } \Omega,$$

then problem (17) has a solution u , with $u(x) \in I$ for every $x \in \overline{\Omega}$.

The proof follows the same lines of the one given in the previous section for the parabolic case. The analogue of Proposition 2 has been first proved by Amann [2] in the case of classical solutions, and can be found in [15] for our setting, while the analogue of Proposition 1 is proved in [34, Lemma 5.10].

The remarks and the corollaries at the end of Section 2 clearly hold in this situation, as well.

Corollary 6 *Given $p > 1$ and a function $\zeta \in L^\infty(\Omega)$, with $\text{ess inf } \zeta > 0$, the problem*

$$\begin{cases} -\Delta u = \zeta(x)|u|^{p-1}u + e(x) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \quad (18)$$

has a solution, provided that $\|e\|_{L^\infty}$ is sufficiently small. More precisely, given $\zeta \in L^\infty(\Omega)$, with $\text{ess inf } \zeta > 0$, for every given $\delta > 0$ there is a $\bar{\Lambda}_\delta > 0$ such that, for every $p > 1$, if

$$\|e\|_{L^\infty} \leq \left(\frac{\bar{\Lambda}_\delta}{(1+\delta)^p} \right)^{\frac{p}{p-1}}, \quad (19)$$

then (18) has a solution.

When $N \geq 3$, problems like (18) are usually considered in the literature under the hypothesis that $p \leq \frac{N+2}{N-2} = 2^* - 1$, where 2^* is the critical exponent for the Sobolev imbedding. See, e.g., the recent paper [12] and the references therein. We do not require such a restriction, provided that $\|e\|_{L^\infty}$ is small.

Corollary 7 *Given $p > 1$ and a function $\zeta \in L^\infty(\Omega)$, with $\text{ess inf } \zeta > 0$, the problem*

$$\begin{cases} -\Delta u = \zeta(x)(u^+)^p + e(x) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \quad (20)$$

has a solution, provided that $e(x) \leq 0$ for a.e. $x \in \Omega$, and $\|e\|_{L^\infty}$ is sufficiently small. More precisely, given $\zeta \in L^\infty(\Omega)$, with $\text{ess inf } \zeta > 0$, for every given $\delta > 0$ there is a $\bar{\Lambda}_\delta > 0$ such that, for every $p > 1$, if $e(x) \leq 0$ for a.e. $x \in \Omega$, and (19) holds, then (20) has a solution.

Problems like (20) have been studied in [36] and [23], assuming, when $N \geq 3$, that $p < \frac{N}{N-2}$. We do not require such a restriction, provided that $\|e\|_{L^\infty}$ is small.

Corollary 8 *Assume that $\zeta_1, \zeta_2 \in L^\infty(\Omega)$ are bounded below by a positive constant, $p > 1$, and $q < p$. Then, there is a $\bar{\lambda} > 0$ such that, for every $\lambda \in [0, \bar{\lambda}]$, problem*

$$\begin{cases} -\Delta u = \zeta_1(x)u^p - \lambda\zeta_2(x)u^q & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega, \end{cases} \quad (21)$$

has a solution.

Similar problems were considered, e.g., in [28] when $N \geq 3$ and $p < 2^* - 1$, and in [35, 13] for $p = 2^* - 1$. If $1 < q < p$, problem (21) has a concave-convex nonlinearity and reminds a similar Dirichlet problem introduced in [5] (see also [10]). However, differently from our situation, lower and upper solutions are found there with the usual ordering $\alpha \leq \beta$. Using this fact, multiplicity results are proved by the use of variational methods, provided that $p \leq 2^* - 1$ when $N \geq 3$. We do not know whether in our case multiplicity results could be obtained.

4 Second order ODE's

In this section, we consider either the Neumann problem

$$\begin{cases} -u'' + a(x)u' = f(x, u, u'), \\ u'(0) = u'(1) = 0, \end{cases} \quad (22)$$

or the periodic problem

$$\begin{cases} -u'' + a(x)u' = f(x, u, u'), \\ u(0) = u(1), \quad u'(0) = u'(1). \end{cases} \quad (23)$$

Here, $a \in L^\infty(0, 1)$, and the function $f : [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is L^1 -Carathéodory. We say that u is a *solution* of problem (22), or (23), if u belongs to $W^{2,1}(0, 1)$, it satisfies the differential equation a.e. in $]0, 1[$ and the boundary conditions pointwise.

Let us define the class of functions

$$\mathcal{F}(I, \Lambda, K),$$

where $I \subseteq \mathbb{R}$ is an interval, and Λ, K are some nonnegative constants. Its elements are the Carathéodory functions $f : [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ which satisfy the following Bernstein-Nagumo growth condition:

$$(BN) \quad \begin{aligned} &|f(x, u, v)| \leq h(x) + K|v|^2, \\ &\text{for a.e. } x \in]0, 1[\text{ and every } (u, v) \in I \times \mathbb{R}, \end{aligned}$$

for some $h \in L^1(0, 1)$, with $\|h\|_{L^1} \leq \Lambda$.

Theorem 5 *Let $I \subseteq \mathbb{R}$ be an interval. Given a compact interval J , contained in the interior of I , there is a $\Lambda > 0$ such that, for every $f \in \mathcal{F}(I, \Lambda, \Lambda)$, if there are two constants α, β in J for which*

$$f(x, \beta, 0) \leq 0 \leq f(x, \alpha, 0) \quad \text{a.e. in }]0, 1[,$$

then problems (22) and (23) have a solution u , with $u(x) \in I$ for every $x \in [0, 1]$.

The proof of the theorem uses the same arguments as in the parabolic case, with some small modifications, due to the fact that, now, $r = N = 1$. The analogue of Proposition 2 is rather standard in this setting (see, e.g., [14], with different kinds of Bernstein-Nagumo conditions). The analogue of Proposition 1 is reported in the Appendix, and provides a uniform estimate in the $W^{2,1}$ -norm. Some more care has to be used in the choice of the spaces, since $W^{2,1}(0, 1)$ is continuously imbedded in $C^1([0, 1])$, but not compactly imbedded. Taking advantage of (BN), one can choose to define the operators \tilde{N}_m on the space $W^{1,2}(0, 1)$ instead of $C^1([0, 1])$, taking values in $L^1(0, 1)$. Since \tilde{N}_m maps bounded sets into bounded sets and

$$W^{2,1}(0, 1) \subset\subset W^{1,2}(0, 1) \subset\subset C([0, 1]),$$

the corresponding operators $\tilde{\mathcal{S}}_m : W^{1,2}(0, 1) \rightarrow W^{1,2}(0, 1)$, defined by (2), are now completely continuous, and the remaining part of the proof is analogous to that of the parabolic case.

The remarks and the corollaries at the end of Section 2 clearly hold in this situation, as well. Theorem 5 generalizes a result by Pournaki and Razani [30, 31], where the existence of periodic solutions with small period has been proved. A simple change of the time variable permits to fix the period and reduce to our case.

Remark 5 Assuming f to be continuous, it is possible to introduce variants of the Bernstein-Nagumo condition, involving only one-sided growth restrictions. For example, instead of $\mathcal{F}(I, \Lambda, K)$, we can introduce the class of functions

$$\mathcal{G}(I, \Lambda, \varphi),$$

where $I \subseteq \mathbb{R}$ is an interval, Λ is a nonnegative constant, and $\varphi : [0, +\infty[\rightarrow]0, +\infty[$ is a continuous function such that

$$\int_1^{+\infty} \frac{s}{\varphi(s)} ds = +\infty.$$

Its elements are the continuous functions $f : [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ which satisfy

$$f(x, u, v) \leq \Lambda \varphi(|v|),$$

for every $(x, u, v) \in [0, 1] \times I \times \mathbb{R}$. For this and other types of one-sided conditions, and related topics, we refer to the classical literature, cf. [20] and the references therein. A comprehensive review can be found in [14].

Remark 6 The left hand side of the differential equation may also contain some type of nonlinear lower order terms. For example, we can deal with a Liénard type equation

$$-u'' + \eta_1(u)u' = f(x, u, u'),$$

or a Rayleigh type equation

$$-u'' + \eta_2(u') = f(x, u, u'),$$

with $\eta_1, \eta_2 : \mathbb{R} \rightarrow \mathbb{R}$ continuous, provided that η_2 has at most a quadratic growth and $\eta_2(0) = 0$. (We did not write the term $a(x)u'$ for simplicity.) Indeed, if $f = 0$, the only solutions of the Neumann or the periodic problems associated to these equations are the constants, as can be seen by multiplying in the first equation by u , in the second one by u'' , and integrating. This fact permits to show, as in the proof of Theorem 3, that the solutions have a small oscillation, when $f \in \mathcal{F}(I, \Lambda, \Lambda)$, or $f \in \mathcal{G}(I, \Lambda, \varphi)$, with Λ sufficiently small.

5 The p -Laplacian

In this section, we consider the Neumann problem

$$\begin{cases} -\Delta_p u = f(x, u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (24)$$

Here, Ω is a bounded domain in \mathbb{R}^N with a C^2 -boundary, Δ_p is the p -Laplacian, $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$, with $p > 1$, and the function $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is L^∞ -Carathéodory.

In this case, the function f does not depend on the gradient of u , so that problem (24) has a variational structure. We consider a weaker notion of solution than in the previous sections. A function u is a *solution* of problem (24) if $u \in W^{1,p}(\Omega) \cap L^\infty(\Omega)$, and

$$\int_\Omega |\nabla u|^{p-2} \nabla u \nabla w = \int_\Omega f(x, u) w,$$

for every $w \in W^{1,p}(\Omega)$. Known regularity results (see, e.g., [25]) imply that such a solution is in $C^{1,\sigma}(\overline{\Omega})$ for some $\sigma > 0$.

Theorem 6 *Let $I \subseteq \mathbb{R}$ be an interval. Given a compact interval J , contained in the interior of I , there is a $\Lambda > 0$ such that, for every function f satisfying*

$$|f(x, u)| \leq \Lambda \quad \text{for a.e. } x \in \Omega \text{ and every } u \in I,$$

if there are two constants α, β in J for which

$$f(x, \beta) \leq 0 \leq f(x, \alpha) \quad \text{a.e. in } \Omega, \quad (25)$$

then problem (24) has a solution u , with $u(x) \in I$ for every $x \in \overline{\Omega}$.

Proof Denote by $\mathcal{R} : L^\infty(\Omega) \rightarrow C(\overline{\Omega})$ the completely continuous operator which associates to any $h \in L^\infty(\Omega)$ the unique solution v of the problem

$$\begin{cases} -\Delta_p v + |v|^{p-2}v = h & \text{in } \Omega, \\ \partial_\nu v = 0 & \text{on } \partial\Omega. \end{cases}$$

We see that (24) is equivalent to the fixed point problem

$$u = \mathcal{T}(u),$$

where $\mathcal{T} : C(\overline{\Omega}) \rightarrow C(\overline{\Omega})$ is the completely continuous operator defined by

$$\mathcal{T}(u) = \mathcal{R}(f(\cdot, u) + |u|^{p-2}u).$$

First of all we consider the situation when f and α, β satisfy (25), with $\alpha \leq \beta$. In this case, we will need no restriction on Λ .

If $\alpha = \beta$ we obviously have a constant solution. If $\alpha < \beta$, define

$$\gamma(u) = \begin{cases} \alpha & \text{if } u \leq \alpha, \\ u & \text{if } \alpha \leq u \leq \beta, \\ \beta & \text{if } u \geq \beta, \end{cases}$$

and consider the modified problem

$$\begin{cases} -\Delta_p u + |u|^{p-2}u = f(x, \gamma(u)) + |\gamma(u)|^{p-2}\gamma(u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (26)$$

We see that the modified problem (26) is equivalent to the fixed point problem

$$u = \widehat{\mathcal{T}}(u),$$

where $\widehat{\mathcal{T}} : C(\overline{\Omega}) \rightarrow C(\overline{\Omega})$ is the completely continuous operator defined by

$$\widehat{\mathcal{T}}(u) = \mathcal{R}(f(\cdot, \gamma(u)) + |\gamma(u)|^{p-2}\gamma(u)).$$

Since the image of $\widehat{\mathcal{T}}$ is contained in a ball B_R , by Schauder's Theorem, $\widehat{\mathcal{T}}$ has a fixed point in B_R , and

$$d_{LS}(I - \widehat{\mathcal{T}}, B_R) = 1.$$

Moreover, if u is any solution of (26), taking $(\alpha - u)^+$ and $(u - \beta)^+$ as test functions in (26) we see that $\alpha \leq u \leq \beta$. Hence u actually solves (24), thus concluding the proof in the case $\alpha < \beta$.

Notice that, in this setting, the pair (α, β) is said to be *admissible* for (24) if $\alpha < \beta$ and any solution u of (24) satisfying $\alpha \leq u \leq \beta$ is such that $\alpha < u < \beta$. Define

$$\mathcal{U}_{(\alpha, \beta)} = \{u \in C(\overline{\Omega}) : \alpha < u < \beta\},$$

which is bounded and open in $C(\overline{\Omega})$. Since \mathcal{T} and $\widehat{\mathcal{T}}$ coincide on $\overline{\mathcal{U}}_{(\alpha, \beta)}$, by the excision property of the degree, we have that, if (α, β) is admissible,

$$d_{LS}(I - \mathcal{T}, \mathcal{U}_{(\alpha, \beta)}) = d_{LS}(I - \widehat{\mathcal{T}}, \mathcal{U}_{(\alpha, \beta)}) = d_{LS}(I - \widehat{\mathcal{T}}, B_R) = 1.$$

As for Theorem 3, we will now provide the proof by contradiction, by taking $\Lambda = \frac{1}{m}$, with $m \geq 1$ a sufficiently large integer. Let $\delta > 0$ be such that $[\min J - \delta, \max J + \delta] \subseteq I$, and fix $\varepsilon \in]0, \frac{\delta}{2}[$. By contradiction, assume that there exist a function f_m such that

$$|f_m(x, u)| \leq \frac{1}{m} \quad \text{for a.e. } x \in \Omega \text{ and every } u \in I,$$

and two constants $\alpha_m, \beta_m \in J$ satisfying

$$f_m(x, \beta_m) \leq 0 \leq f_m(x, \alpha_m) \quad \text{a.e. in } \Omega,$$

for which the problem

$$\begin{cases} -\Delta_p u = f_m(x, u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (27)$$

has no solution u , with $u(x) \in I$ for every $x \in \overline{\Omega}$. By the above, it has to be $\beta_m < \alpha_m$. Define the modified function

$$\tilde{f}_m(x, u) = \begin{cases} \frac{1}{m} & \text{if } u \leq \beta_m - 2\varepsilon, \\ \frac{1}{m} + \frac{1}{\varepsilon}(f_m(x, u) - \frac{1}{m})(u - \beta_m + 2\varepsilon) & \text{if } \beta_m - 2\varepsilon \leq u \leq \beta_m - \varepsilon, \\ f_m(x, u) & \text{if } \beta_m - \varepsilon \leq u \leq \alpha_m + \varepsilon, \\ -\frac{1}{m} + \frac{1}{\varepsilon}(f_m(x, u) + \frac{1}{m})(\alpha_m + 2\varepsilon - u) & \text{if } \alpha_m + \varepsilon \leq u \leq \alpha_m + 2\varepsilon, \\ -\frac{1}{m} & \text{if } u \geq \alpha_m + 2\varepsilon, \end{cases}$$

and consider the problem

$$\begin{cases} -\Delta_p u = \tilde{f}_m(x, u) & \text{in } \Omega, \\ \partial_\nu u = 0 & \text{on } \partial\Omega. \end{cases} \quad (28)$$

Notice that, as $\Lambda = \frac{1}{m}$, we have $|\tilde{f}_m(x, u)| \leq \frac{1}{m}$ for every $u \in \mathbb{R}$, and

$$\tilde{f}_m(x, \alpha_m + \delta) < 0 < \tilde{f}_m(x, \beta_m - \delta) \quad \text{a.e. in } \Omega.$$

If u is a solution of (28), taking $(\beta_m - 2\varepsilon - u)^+$ as test function we see that $u \geq \beta_m - 2\varepsilon$, so that $u > \beta_m - \delta$. Similarly, taking $(u - \alpha_m - 2\varepsilon)^+$ as test function, one has $u < \alpha_m + \delta$. So, the pair $(\beta_m - \delta, \alpha_m + \delta)$ is admissible for (28).

Using the above information on the degree, we now proceed exactly as in Section 2 to prove the existence of a solution u_m of (28) such that

$$\beta_m - \delta < u_m < \alpha_m + \delta, \quad (29)$$

and, for some $x_m \in \overline{\Omega}$,

$$\beta_m \leq u_m(x_m) \leq \alpha_m. \quad (30)$$

(See (3) and (4).) We can now write $u_m = \mathcal{R}(h_m)$, where

$$h_m(x) = \tilde{f}_m(x, u_m(x)) + |u_m(x)|^{p-2}u_m(x).$$

By (29), there is a constant $c > 0$ such that

$$\|h_m\|_{L^\infty} \leq c, \quad \text{for every } m.$$

Since \mathcal{R} is completely continuous, there are a subsequence, still denoted by $(u_m)_m$, and a function $\bar{u} \in C(\overline{\Omega})$ such that $u_m \rightarrow \bar{u}$ uniformly in $\overline{\Omega}$. As $|\tilde{f}_m(x, u_m(x))| \leq \frac{1}{m}$, for a.e. $x \in \Omega$, it has to be

$$\|h_m - |\bar{u}|^{p-2}\bar{u}\|_{L^\infty} \rightarrow 0,$$

so that $\bar{u} = \mathcal{R}(|\bar{u}|^{p-2}\bar{u})$, i.e.

$$\begin{cases} -\Delta_p \bar{u} = 0 & \text{in } \Omega, \\ \partial_\nu \bar{u} = 0 & \text{on } \partial\Omega. \end{cases}$$

Hence, \bar{u} is constant. Since $u_m \rightarrow \bar{u}$ uniformly in $\overline{\Omega}$, for m sufficiently large we have that $\text{osc } u_m \leq \varepsilon$, and hence, by (30),

$$\beta_m - \varepsilon \leq u_m \leq \alpha_m + \varepsilon.$$

So, u_m is a solution of (P_m) , and $u_m(x) \in I$, for every $x \in \overline{\Omega}$, a contradiction. ■

To conclude, one can easily verify that the remarks and the corollaries in Sections 2 and 3 can be adapted to this situation, as well.

6 Appendix

For the reader's convenience we now give the proof of the following analogue of Proposition 1, which was needed in Section 4 in the framework of ordinary differential equations.

Proposition 3 *Given $M, \Lambda, K > 0$, there is a constant $C > 0$ such that, if $u \in W^{2,1}(0, 1)$ verifies*

$$\begin{cases} |u''| \leq h(x) + K|u'|^2, & \text{a.e. in }]0, 1[, \\ |u| \leq M & \text{in } [0, 1], \end{cases} \quad (31)$$

for some $h \in L^1(0, 1)$ with $\|h\|_{L^1} \leq \Lambda$, then

$$\|u\|_{W^{2,1}} \leq C.$$

Proof Taking if necessary the positive part of h , we can assume that $h \geq 0$ a.e. in $]0, 1[$. We first consider the case when $h \in L^\infty(0, 1)$ and define

$$\varphi(s) = \|h\|_{L^\infty} + 1 + Ks^2.$$

Since $-M \leq u \leq M$, by Lagrange's Theorem, we can find $\xi \in]0, 1[$ such that $|u'(\xi)| \leq 2M$. If $\|u'\|_{L^\infty} \leq 2M$, using (31), we have

$$\|u\|_{W^{2,\infty}} = \|u\|_{L^\infty} + \|u'\|_{L^\infty} + \|u''\|_{L^\infty} \leq 3M + (\|h\|_{L^\infty} + 4KM^2).$$

Otherwise, let $[x_1, x_2] \subseteq [0, 1]$ be an interval such that

$$|u'(x)| > 2M \quad \text{for every } x \in]x_1, x_2[,$$

and, either $|u'(x_1)| = 2M$, or $|u'(x_2)| = 2M$. Let $i = 1, 2$ be the index for which $|u'(x_i)| = 2M$. Then, considering separately the cases when u' is positive or negative on $[x_1, x_2]$, one easily verifies that for every $x \in [x_1, x_2]$,

$$\int_{2M}^{|u'(x)|} \frac{s}{\varphi(s)} ds \leq \left| \int_{x_i}^x \frac{|u'(t)| |u''(t)|}{\varphi(|u'(t)|)} dt \right| \leq \left| \int_{x_i}^x |u'(t)| dt \right| = |u(x) - u(x_i)| \leq 2M.$$

Since $\int_0^{+\infty} \frac{s}{\varphi(s)} ds = +\infty$, there is a constant $R > 0$, depending only on $\|h\|_{L^\infty}$, K , and M , such that

$$|u'(x)| \leq R \quad \text{for every } x \in [x_1, x_2].$$

This proves that $\|u'\|_{L^\infty} \leq R$. Using (31), we then have

$$\|u\|_{W^{2,\infty}} \leq M + R + (\|h\|_{L^\infty} + KR^2),$$

and the proof is completed in this case.

Assume now $h \in L^1(0, 1)$ and, following [34], set

$$e(x) = \frac{-u''(x) + u(x)}{1 + h(x) + K|u'(x)|^2}.$$

Then, being $h \geq 0$ a.e. in $]0, 1[$, it is $e \in L^\infty(0, 1)$, with $\|e\|_{L^\infty} \leq 1 + M$.

Let $v \in W^{2,1}(0, 1)$ be the solution of

$$\begin{cases} -v'' + v = e(x)(1 + h(x)), \\ v(0) = v(1), \quad v'(0) = v'(1). \end{cases}$$

Then, there is a constant $\bar{c} > 0$ such that

$$\|v\|_{C^1} \leq \bar{c}\|e(1 + h)\|_{L^1} \leq \bar{c}(1 + M)(1 + \|h\|_{L^1}).$$

Let now $w = u - v$. Then

$$-w'' + w = Ke(x)|u'(x)|^2,$$

so that, by (31),

$$\begin{aligned} |w''(x)| &\leq |w(x)| + K|e(x)||u'(x)|^2 \\ &\leq |u(x)| + |v(x)| + K|e(x)|(2|v'(x)|^2 + 2|w'(x)|^2) \\ &\leq C_1 + C_2|w'(x)|^2, \end{aligned}$$

for a.e. $x \in]0, 1[$, where

$$C_1 = M + \bar{c}(1 + M)(1 + \|h\|_{L^1}) + 2K(1 + M)(\bar{c}(1 + M)(1 + \|h\|_{L^1}))^2$$

and $C_2 = 2K(1 + M)$. Moreover,

$$|w(x)| \leq |u(x)| + |v(x)| \leq M + \bar{c}(1 + M)(1 + \|h\|_{L^1}),$$

for every $x \in [0, 1]$.

Therefore, w verifies the same assumptions as u with a constant function $h = C_1$, and, by the first part of the proof, there is a constant $R > 0$ depending only on M , C_1 , and C_2 , such that $\|w'\|_{L^\infty} \leq R$. Then,

$$\|u'\|_{L^\infty} \leq \|w'\|_{L^\infty} + \|v'\|_{L^\infty} \leq R + \bar{c}(1 + M)(1 + \|h\|_{L^1}).$$

By (31), we have

$$\|u''\|_{L^1} \leq \|h\|_{L^1} + K[R + \bar{c}(1 + M)(1 + \|h\|_{L^1})]^2,$$

which concludes the proof. ■

Acknowledgement. We wish to thank Pierpaolo Omari and Fabio Zanolin for their fruitful comments and suggestions.

References

- [1] Adimurti and G. Mancini, The Neumann problem for elliptic equations with critical non-linearity, in: *Nonlinear Analysis*, Scuola Normale Sup. Pisa, pp. 9–25, 1991.
- [2] H. Amann, On the number of solutions of nonlinear equations in ordered Banach spaces, *J. Funct. Anal.* 11 (1972), 346–384.
- [3] H. Amann, Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces, *SIAM Review* 18 (1976), 620–709.
- [4] H. Amann, A. Ambrosetti and G. Mancini, Elliptic equations with non-invertible Fredholm linear part and bounded nonlinearities, *Math. Z.* 158 (1978), 179–194.
- [5] A. Ambrosetti, H. Brezis and G. Cerami, Combined effects of concave and convex nonlinearities in some elliptic problems, *J. Funct. Anal.* 122 (1994) 519–543.
- [6] A. Bahri, Topological results on a certain class of functionals and applications, *J. Funct. Anal.* 41 (1981), 397–427.
- [7] A. Bahri and H. Berestycki, A perturbation method in critical point theory and applications, *Trans. Amer. Math. Soc.* 267 (1981), 1–32.
- [8] A. Bahri and P.-L. Lions, Morse index of some min-max critical points. I. Application to multiplicity results, *Comm. Pure Appl. Math.* 41 (1988), 1027–1037.
- [9] T. Bartsch and M. Willem, On an elliptic equation with concave and convex nonlinearities, *Proc. Amer. Math. Soc.* 123 (1995), 3555–3561.
- [10] L. Boccardo, M. Escobedo and I. Peral, A Dirichlet problem involving critical exponents, *Nonlinear Anal.* 24 (1995), 1639–1648.
- [11] H. Brezis and L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.* 36 (1983), 436–477.
- [12] J. Chabrowski, The critical Neumann problem for semilinear elliptic equations with concave perturbations, *Ricerche Mat.* 56 (2007), 297–319.
- [13] J. Chabrowski and M. Willem, Least energy solutions of a critical Neumann problem with a weight, *Calc. Var.* 15 (2002), 421–431.
- [14] C. De Coster and P. Habets, *Two-Point Boundary Value Problems, Lower and Upper Solutions*, Elsevier, Amsterdam, 2006.

- [15] C. De Coster and M. Henrard, Existence and localization of solution for elliptic problem in presence of lower and upper solutions without any order, *J. Differential Equations* 145 (1998), 420–452.
- [16] C. De Coster, F. Obersnel and P. Omari, A qualitative analysis, via lower and upper solutions, of first order periodic evolutionary equations with lack of uniqueness, in: *Handbook of Differential Equations, ODE's*, A. Canada, P. Drabek and A. Fonda Eds., Elsevier, Amsterdam, 2006.
- [17] C. De Coster and P. Omari, Unstable periodic solutions of a parabolic problem in presence of non-well-ordered lower and upper solutions, *J. Funct. Anal.* 175 (2000), 52–88.
- [18] C. De Coster and P. Omari, Stability and instability in periodic parabolic problems via lower and upper solutions, *Quaderni Matematici* 539 (2003), Univ. Trieste.
- [19] M. Furi and M.P. Pera, Co-bifurcating branches of solutions for nonlinear eigenvalue problems in Banach spaces, *Ann. Mat. Pura Appl.* 135 (1983), 119–132.
- [20] R.E. Gaines and J. Mawhin, *Coincidence Degree and Nonlinear Differential Equations*, *Lect. Notes Math.* 568, Springer, Berlin, 1977.
- [21] J.-P. Gossez and P. Omari, A necessary and sufficient condition of nonresonance for a semilinear Neumann problem, *Proc. Amer. Math. Soc.* 114 (1992), 433–442.
- [22] P. Habets and P. Omari, Existence and localization of solutions of second order elliptic problems using lower and upper solutions in the reversed order, *Top. Meth. Nonlinear Anal.* 8 (1996), 25–56.
- [23] R. Kannan and R. Ortega, Landesman-Lazer conditions for problems with “one-side bounded” nonlinearities, *Nonlinear Anal.* 9 (1985), 1313–1317.
- [24] S. Li and Z. Liu, Multiplicity of solutions for some elliptic equations involving critical and supercritical Sobolev exponents. *Topol. Methods Nonlinear Anal.* 28 (2006), 235–261.
- [25] G.M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations, *Nonlinear Anal.* 12 (1988), 1203–1219.
- [26] G.M. Lieberman, Time-periodic solutions of linear parabolic differential equations, *Comm. P.D.E.* 24 (1999), 631–663.
- [27] J. Mawhin and K. Schmitt, Upper and lower solutions and semilinear second order elliptic equations with non-linear boundary conditions, *Proc. Royal Soc. Edinburgh* 97A (1984), 199–207.

- [28] W.M. Ni and I. Takagi, On the Neumann problem for some semilinear elliptic equations and systems of activator-inhibitor type, *Trans. Amer. Math. Soc.* 297 (1986) 351–368.
- [29] P. Omari, Non-ordered lower and upper solutions and solvability of the periodic problem for the Liénard and the Rayleigh equations, *Rend. Ist. Mat. Univ. Trieste* 20 (1988), 54–64.
- [30] M.R. Pournaki and A. Razani, On the existence of periodic solutions for a class of generalized forced Liénard equations, *Appl. Math. Letters* 20 (2007) 248–254.
- [31] M.R. Pournaki and A. Razani, Erratum to “On the existence of periodic solutions for a class of generalized forced Liénard equations”, *Appl. Math. Letters*, to appear.
- [32] P.H. Rabinowitz, Multiple critical points of perturbed symmetric functionals, *Trans. Amer. Math. Soc.* 272 (1982), 753–770.
- [33] M. Struwe, Infinitely many critical points for functionals which are not even and applications to superlinear boundary value problems, *Manuscripta Math.* 32 (1980), 335–364.
- [34] G.M. Troianiello, *Elliptic differential equations and obstacle problems*, Plenum Press, New York, 1987.
- [35] X.-J. Wang, Neumann problems for semilinear elliptic equations involving critical Sobolev exponents, *J. Differential Equations* 93 (1991), 283–310.
- [36] J.R. Ward, Perturbations with some superlinear growth for a class of second order elliptic boundary value problems, *Nonlinear Anal.* 6 (1982), 367–374.